

**NSF–CBMS CONFERENCE ON  
THE CAHN–HILLIARD EQUATION:  
RECENT ADVANCES AND APPLICATIONS  
PRELIMINARY READING**

ALAIN MIRANVILLE

1. THE CAHN–HILLIARD EQUATION

The Cahn–Hilliard system

$$(1.1) \quad \frac{\partial u}{\partial t} = \kappa \Delta \mu, \quad \kappa > 0, \quad \mu = -\alpha \Delta u + f(u), \quad \alpha > 0,$$

usually rewritten, equivalently, as the fourth-order in space parabolic equation

$$(1.2) \quad \frac{\partial u}{\partial t} + \alpha \kappa \Delta^2 u - \kappa \Delta f(u) = 0$$

which is precisely the equation known as the Cahn–Hilliard equation, was proposed by J.W. Cahn<sup>1</sup> and J.E. Hilliard in 1958 (see [63]). These equations play an essential role in materials science as they describe important qualitative features of two-phase systems related with phase separation processes, assuming isotropy and a constant temperature. This can be observed, e.g., when a binary alloy (e.g., Aluminium/Zinc, see [390], or Iron/Chromium, see [309, 310, 311]) is cooled down sufficiently. One then observes a partial nucleation (i.e., the apparition of nuclides in the material) or a total nucleation, known as spinodal decomposition: the initially homogeneous material quickly becomes inhomogeneous, resulting in a very finely dispersed microstructure. In a second stage, which is called coarsening and occurs at a slower time scale, these microstructures coarsen. Such phenomena play an essential role in the mechanical properties of the material, e.g., strength, hardness, fracture, toughness and ductility. We refer the reader to, e.g., [61, 63, 275, 280, 302, 303, 345, 347] for more details.

Here,  $u$  is the order parameter (we will consider a rescaled density of atoms or concentration of one of the material’s components which takes values between  $-1$

---

<sup>1</sup>John Werner Cahn (January 9, 1928–March 14, 2016) played a major role in materials science.

and 1,  $-1$  and 1 corresponding to the pure states; the density of the second component is  $-u$ , meaning that the total density is a conserved quantity<sup>2</sup>) and  $\mu$  is the chemical potential (more precisely, the difference of chemical potentials between the two components). Furthermore,  $f$  is the derivative of a double-well potential  $F$ . A thermodynamically relevant potential  $F$  is the following logarithmic function which follows from a mean-field model:

$$(1.3) \quad F(s) = \frac{\theta_c}{2}(1-s^2) + \frac{\theta}{2}[(1-s)\ln(\frac{1-s}{2}) + (1+s)\ln(\frac{1+s}{2})], \quad s \in (-1, 1),$$

$$0 < \theta < \theta_c,$$

i.e.,

$$(1.4) \quad f(s) = -\theta_c s + \frac{\theta}{2} \ln \frac{1+s}{1-s},$$

although such a function is very often approximated by regular ones, typically,  $F(s) = \frac{1}{4}(s^2-1)^2$ , i.e.,  $f(s) = s^3 - s$ ; more generally, one can take  $F(s) = \frac{1}{4}(s^2 - \beta^2)^2$ ,  $\beta \in \mathbb{R}$ . The logarithmic terms in (1.3) correspond to the entropy of mixing and  $\theta$  and  $\theta_c$  are proportional to the absolute temperature (assumed constant during the process) and a critical temperature, respectively; the condition  $\theta < \theta_c$  ensures that  $F$  has indeed a double-well form and that phase separation can occur. Also note that the polynomial approximation is reasonable when the quench is shallow, i.e., when the absolute temperature is close to the critical one. Finally,  $\kappa$  is the mobility and  $\alpha$  is related to the surface tension at the interface.

From a phenomenological point of view, the Cahn–Hilliard system can be derived as follows.

One considers the following (total) free energy, called Ginzburg–Landau free energy:

$$(1.5) \quad \Psi_\Omega(u, \nabla u) = \int_\Omega \left( \frac{\alpha}{2} |\nabla u|^2 + F(u) \right) dx,$$

where  $\Omega \subset \mathbb{R}^n$ ,  $n = 1, 2$  or  $3$ , is the domain occupied by the material. The gradient term in (1.5) was proposed in [63] in order to model the surface energy of the interface (i.e., capillarity; note that such gradients go back to J.D. van der Waals, see [395]);  $F$  is also called homogeneous free energy.

One then has the mass balance

---

<sup>2</sup>If  $u_A$  and  $u_B$  denote the densities of the two components, then, before rescaling, one has  $u_A + u_B = 1$ . Replacing  $u$  by  $2u - 1$ , one obtains, after rescaling,  $u_A + u_B = 0$ .

$$(1.6) \quad \frac{\partial u}{\partial t} = -\operatorname{div} h,$$

where  $h$  is the mass flux which is related to the chemical potential  $\mu$  by the following (postulated) constitutive equation which resembles the Fick's law:

$$(1.7) \quad h = -\kappa \nabla \mu.$$

The usual definition of the chemical potential is that it is the derivative of the free energy with respect to the order parameter. Here, such a definition is incompatible with the presence of  $\nabla u$  in the free energy. Instead,  $\mu$  is defined as a variational derivative of the free energy with respect to  $u$ , which yields (assuming proper boundary conditions)

$$(1.8) \quad \mu = -\alpha \Delta u + f(u);$$

hence the Cahn–Hilliard system. This variational derivative can be (formally) seen by writing that, for a small variation,

$$\delta \Psi_{\Omega} = \int_{\Omega} (\alpha \nabla u \cdot \nabla \delta u + f(u) \delta u) dx,$$

where  $\cdot$  denotes the usual Euclidean scalar product. Assuming compatible boundary conditions and integrating by parts, this yields

$$\delta \Psi_{\Omega} = \int_{\Omega} (-\alpha \Delta u + f(u)) \delta u dx,$$

from which the definition follows.

The Cahn–Hilliard system, in a bounded and regular domain  $\Omega$ , usually is endowed with Neumann boundary conditions, namely,

$$(1.9) \quad \frac{\partial \mu}{\partial \nu} = 0, \text{ on } \Gamma,$$

meaning that there is no mass flux at the boundary (note that  $h \cdot \nu = -\kappa \frac{\partial \mu}{\partial \nu}$ ), and

$$(1.10) \quad \frac{\partial u}{\partial \nu} = 0, \text{ on } \Gamma,$$

which is a natural variational boundary condition (by natural, we mean that it allows to write down a convenient variational/weak formulation in view of the mathematical analysis of the problem; this boundary condition also yields that the interface is orthogonal to the boundary). Here,  $\Gamma = \partial \Omega$  and  $\nu$  is the unit outer normal to the

boundary. In particular, it follows from the first boundary condition that we have the conservation of mass, i.e., of the spatial average of the order parameter, obtained by (formally) integrating the first equation of (1.1) over  $\Omega$ ,

$$(1.11) \quad \langle u(t) \rangle \equiv \frac{1}{\text{Vol}(\Omega)} \int_{\Omega} u(t, x) dx = \langle u(0) \rangle, \quad \forall t \geq 0.$$

If we have in mind the fourth-order in space Cahn–Hilliard equation, we can rewrite these boundary conditions, equivalently, as

$$(1.12) \quad \frac{\partial u}{\partial \nu} = \frac{\partial \Delta u}{\partial \nu} = 0, \quad \text{on } \Gamma.$$

We can also consider periodic boundary conditions (in which case  $\Omega = \Pi_{i=1}^n(0, L_i)$ ,  $L_i > 0$ ,  $i = 1, \dots, n$ ); in that case, we still have the conservation of mass. Note that we generally do not consider Dirichlet boundary conditions, due precisely to the fact that they do not yield the conservation of mass, although such boundary conditions certainly simplify the mathematical analysis.

Now, the question of how the phase separation process (i.e., the spinodal decomposition) is influenced by the presence of walls has gained much attention (see [164, 165, 265] and the references therein). This problem has mainly been studied for polymer mixtures (although it should also be important for other systems, such as binary metallic alloys): from a technological point of view, binary polymer mixtures are particularly interesting, since the occurring structures during the phase separation process may be frozen by a rapid quench into the glassy state; microstructures at surfaces on very small length scales can be produced in this way.

We also recall that the usual variational boundary condition  $\frac{\partial u}{\partial \nu} = 0$  on the boundary yields that the interface is orthogonal to the boundary, meaning that the contact line, when the interface between the two components meets the walls, is static, which is not reasonable in many situations. This is the case, e.g., for mixtures of two immiscible fluids: in that case, the contact angle should be dynamic, due to the movements of the fluids. This can also be the case in the context of binary alloys, whence the need to define dynamic boundary conditions for the Cahn–Hilliard equation.

In that case, we again write that there is no mass flux at the boundary (i.e., that (1.9) still holds). Then, in order to obtain the second boundary condition, following the phenomenological derivation of the Cahn–Hilliard system, we consider, in addition to the usual Ginzburg–Landau free energy and assuming that the interactions with the walls are short-ranged, a surface free energy of the form

$$(1.13) \quad \Psi_{\Gamma}(u, \nabla_{\Gamma} u) = \int_{\Gamma} \left( \frac{\alpha_{\Gamma}}{2} |\nabla_{\Gamma} u|^2 + G(u) \right) d\sigma, \quad \alpha_{\Gamma} > 0,$$

where  $\nabla_\Gamma$  is the surface gradient and  $G$  is a surface potential. Thus, the total free energy of the system reads

$$(1.14) \quad \Psi = \Psi_\Omega + \Psi_\Gamma.$$

Writing finally that the system tends to minimize the excess surface energy, we are led to postulate the following boundary condition:

$$(1.15) \quad \frac{1}{d} \frac{\partial u}{\partial t} - \alpha_\Gamma \Delta_\Gamma u + g(u) + \alpha \frac{\partial u}{\partial \nu} = 0, \text{ on } \Gamma,$$

i.e., there is a relaxation dynamics on the boundary. This boundary condition usually is referred to as dynamic boundary condition, in the sense that the kinetics, i.e.,  $\frac{\partial u}{\partial t}$ , appears explicitly. Here,  $\Delta_\Gamma$  is the Laplace–Beltrami operator,  $g = G'$  and  $d > 0$  is some relaxation parameter. Furthermore, in the original derivation, one has  $G(s) = \frac{1}{2}a_\Gamma s^2 - b_\Gamma s$ , where  $a_\Gamma > 0$  accounts for a modification of the effective interaction between the components at the walls and  $b_\Gamma$  characterizes the possible preferential attraction (or repulsion) of one of the components by the walls (when  $b_\Gamma$  vanishes, there is no preferential attraction). We can note that it follows from the boundary conditions that, formally,

$$\frac{d\Psi}{dt} = -\frac{1}{\kappa} \left\| \frac{\partial u}{\partial t} \right\|_{H^{-1}(\Omega)}^2 - \frac{1}{d} \left\| \frac{\partial u}{\partial t} \right\|_{L^2(\Gamma)}^2 \leq 0,$$

where  $\|\cdot\|_X$  denotes the norm on the Banach space  $X$ ; in the case of the classical Neumann boundary conditions, one has

$$\frac{d\Psi_\Omega}{dt} = -\frac{1}{\kappa} \left\| \frac{\partial u}{\partial t} \right\|_{H^{-1}(\Omega)}^2 \leq 0.$$

We also refer the reader to [30, 166] for other physical derivations of the dynamic boundary condition, obtained by taking the continuum limit of lattice models within a direct mean-field approximation and by applying a density functional theory, respectively, to [362] for the derivation of dynamic boundary conditions in the context of two-phase fluid flows and to [372, 377] for an approach based on concentrated capacity.

Actually, it would seem more reasonable, in the case of nonpermeable walls, to write the conservation of mass both in the bulk  $\Omega$  and on the boundary  $\Gamma$ , i.e.,

$$\frac{d}{dt} \left( \int_\Omega u \, dx + \int_\Gamma u \, d\sigma \right) = 0.$$

Indeed, due to the interactions with the walls, one should expect some mass on the boundary. We assume that the first equation of (1.1) still holds. Then, writing that

$$\mu = \partial_u \Psi,$$

where  $\partial$  is the variational derivative mentioned above (note that, in the original derivation, one has  $\mu = \partial_u \Psi_\Omega$ ), we obtain the second equation of (1.1), together with the boundary condition

$$\mu = -\alpha_\Gamma \Delta_\Gamma u + g(u) + \alpha \frac{\partial u}{\partial \nu}, \text{ on } \Gamma.$$

We now note that, owing to the first equation of (1.1), the above mass conservation reads

$$\int_\Gamma \left( \frac{\partial u}{\partial t} + \kappa \frac{\partial \mu}{\partial \nu} \right) d\sigma = 0.$$

A class of boundary conditions which ensure this mass conservation reads

$$\frac{\partial u}{\partial t} + \beta_\Gamma \Delta_\Gamma u + \kappa \frac{\partial \mu}{\partial \nu} = 0, \text{ on } \Gamma, \beta_\Gamma \geq 0.$$

We can thus see that, when  $\beta_\Gamma > 0$ , we also have a Cahn–Hilliard type system on the boundary. Note that, when  $\beta_\Gamma = 0$ , it follows from the above that

$$\frac{d\Psi}{dt} = -\kappa \|\nabla \mu\|_{L^2(\Omega)^n}^2 \leq 0.$$

Similar dynamic boundary conditions, in the case of semipermeable walls, are considered in [174, 175, 185]. Furthermore, in [294], based on an energetic variational approach and Onsager’s principle of maximum energy dissipation, one recovers these dynamic boundary conditions, together with the no mass flux condition (1.9); in that case, one has mass conservation in the bulk and on the boundary, separately.

The Cahn–Hilliard system/equation is now quite well understood, at least from a mathematical point of view. In particular, one has a rather complete picture as far as the existence, the uniqueness and the regularity of solutions and the asymptotic behavior of the associated dynamical system are concerned. We refer the reader to (among a huge literature), e.g., [5, 35, 59, 83, 89, 93, 96, 109, 122, 131, 141, 143, 147, 149, 152, 174, 175, 185, 198, 206, 207, 212, 264, 278, 287, 294, 297, 300, 324, 333, 334, 337, 341, 342, 343, 344, 345, 347, 360, 363, 370, 388, 412, 424]. As far as the asymptotic behavior of the system is concerned, one has, in particular, the existence of finite-dimensional attractors. Such sets give information on the global/ all possible dynamics of the system. Furthermore, the finite dimensionality means, very roughly speaking, that, even though the initial phase space is infinite-dimensional, the limit dynamics can be described by a finite number of parameters. We refer the interested

reader to, e.g., [19, 97, 134, 335, 388] for more details and discussions on this. One also has the convergence of single trajectories to steady states.

We can note that we assumed so far that the mobility  $\kappa$  is a strictly positive constant. Actually,  $\kappa$  is often expected to depend on the order parameter and to degenerate at the singular points of  $f$  in the case of a logarithmic nonlinear term (see [62, 147, 148, 191, 417]; see also [421] for a discussion in the context of immiscible binary fluids). Note however that this essentially restricts the diffusion process to the interfacial region and is observed, typically, in physical situations in which the movements of atoms are confined to this region (see [364]). In that case, the first equation of (1.1) reads

$$\frac{\partial u}{\partial t} = \operatorname{div}(\kappa(u)\nabla\mu),$$

where, typically,  $\kappa(s) = 1 - s^2$ . In particular, the existence of solutions to the Cahn–Hilliard equation with degenerate mobilities and logarithmic nonlinearities is proved in [147]; note that, up to now, only existence of weak solutions is known, nothing else. The asymptotic behavior, and, more precisely, the existence of attractors, of the Cahn–Hilliard equation with nonconstant and nondegenerating mobilities is studied in [378, 379].

We can also note that the gradient term in the Ginzburg–Landau free energy (1.5) accounts for the fact that the interactions between the material’s components are assumed to be short-ranged. Actually, this term is obtained by approximation of a nonlocal term which also accounts for long-ranged interactions (see [63]). The Cahn–Hilliard equation, with a nonlocal term, was derived rigorously by G. Giacomin and J.L. Lebowitz in [204, 205], based on stochastic arguments, by considering a lattice gas with long range Kac potentials (i.e., the interaction energy between two particles at  $x$  and  $y$  ( $x, y \in \mathcal{Z}^n$ ) is given by  $\gamma^n \mathcal{K}(\gamma|x - y|)$ ,  $\gamma > 0$  being sent to 0 and  $\mathcal{K}$  being a smooth function). In that case, the (total) free energy reads

$$(1.16) \quad \Psi_\Omega(u) = \int_{\mathbb{T}^n} [f(u(x)) + u(x) \int_{\mathbb{T}^n} \mathcal{K}(|x - y|)(1 - u(y)) dy] dx,$$

where  $\mathbb{T}^n$  is the  $n$ -dimensional torus. Furthermore, rewriting the total free energy in the form

$$\Psi_\Omega(u) = \int_{\mathbb{T}^n} [f(u(x)) + k_1(x)u(x)(1 - u(x)) + \frac{1}{2} \int_{\mathbb{T}^n} \mathcal{K}(|x - y|)|u(x) - u(y)|^2 dy] dx,$$

where  $k_1(x) = \int_{\mathbb{T}^n} \mathcal{K}(|x - y|) dy$ , one can, by expanding the last term and keeping only some terms in the expansion, recover the Ginzburg–Landau free energy (this is reasonable when the scale on which the free energy varies is large compared with

$\gamma^{-1}$ ; the macroscopic evolution is observed here on the spatial scale  $\gamma^{-1}$  and time scale  $\gamma^{-2}$ ); see also [307]. Such models were studied, e.g., in [3, 27, 173, 176, 177, 178, 181, 254] (see also [78, 133, 230, 231, 232, 415] for the numerical analysis and simulations).

Now, it is interesting to note that the Cahn–Hilliard equation and some of its variants are also relevant in other phenomena than phase separation in binary alloys. We can mention, for instance, dealloying (this can be observed in corrosion processes, see [154]), population dynamics (see [100]), tumor growth (see [18, 266]), bacterial films (see [273]), thin films (see [349, 391]), chemistry (see [396]), image processing (see [28, 29, 72, 79, 132]) and even in astronomy, with the rings of Saturn (see [394]), and ecology (for instance, the clustering of mussels can be perfectly well described by the Cahn–Hilliard equation, see [296] (see also Youtube,

<https://www.youtube.com/watch?v=u-mEjfBaYks>

and

<https://www.youtube.com/watch?v=OYcXZ7Ho4o8>,

for real mussels clustering and simulations, respectively); of course, in that case the time scale is much larger, typically, weeks or months).

In particular, several such phenomena can be modeled by the following generalized Cahn–Hilliard equation:

$$(1.17) \quad \frac{\partial u}{\partial t} + \alpha \kappa \Delta^2 u - \kappa \Delta f(u) + g(x, u) = 0, \quad \alpha, \kappa > 0$$

(here,  $\alpha$  and  $\kappa$  do not necessarily have the same physical meaning as in the original Cahn–Hilliard equation). The above general equation contains, in particular, the following models:

(i) Mixed Allen–Cahn/Cahn–Hilliard system. In that case, we consider the following system of equations:

$$\frac{\partial u}{\partial t} = \varepsilon^2 D \Delta \mu - \mu, \quad D, \varepsilon > 0, \quad \mu = -\Delta u + \frac{f(u)}{\varepsilon^2},$$

which can be rewritten, equivalently, as

$$\frac{\partial u}{\partial t} + \varepsilon^2 D \Delta^2 u - \Delta(Df(u) + u) + \frac{f(u)}{\varepsilon^2} = 0$$

and is indeed of the form above. In particular, without the term  $\varepsilon^2 D \Delta \mu$  in the first equation, we have the Allen–Cahn equation and, without the term  $-\mu$ , we have the Cahn–Hilliard equation. These equations were proposed in order to account for microscopic mechanisms such as surface diffusion and adsorption/desorption (see [258, 260, 261, 308]) and were studied in [250, 251, 252, 253, 259].



(ii) Cahn–Hilliard–Oono<sup>3</sup> equation (see [317, 348, 397]). In that case,

$$g(x, s) = g(s) = \beta s, \quad \beta > 0.$$

This function was proposed in [348] in order to account for long-ranged (i.e., nonlocal) interactions in phase separation, but also to simplify numerical simulations, due to the fact that we do not have to account for the conservation of mass, although it seems that this equation has never been considered in simulations.

Actually, it can be surprising that nonlocal interactions can be described by such a simple linear term. This can be seen by noting that one considers here the free energy

$$(1.18) \quad \Psi_\Omega = \int_\Omega \left( \frac{\alpha}{2} |\nabla u|^2 + F(u) + \int_\Omega u(y)g(y, x)u(x) dy \right) dx,$$

where the function  $g$  describes the long-ranged interactions. In particular, in Oono’s model and in three space dimensions, one takes

$$(1.19) \quad g(y, x) = \frac{4\pi\beta}{|y - x|}.$$

Note that the long-ranged interactions are repulsive when  $u(y)$  and  $u(x)$  have opposite signs and thus favor the formation of interfaces (see [397] and the references therein). Writing finally, as in the derivation of the classical Cahn–Hilliard equation,

$$(1.20) \quad \frac{\partial u}{\partial t} = \kappa \Delta \partial_u \Psi_\Omega,$$

we find the Cahn–Hilliard–Oono equation, noting that  $-\frac{1}{|y-x|}$  is the Green function associated with the Laplace operator (see [397] and the references therein for more details).

A variant of this model, proposed in [95] to model microphase separation of diblock copolymers, consists in taking

$$g(x, s) = g(s) = \beta \left( s - \frac{1}{\text{Vol}(\Omega)} \int_\Omega u_0(x) dx \right), \quad \beta > 0,$$

where  $u_0$  is the initial condition. In that case, we have the conservation of mass and efficient simulations, based on multigrid solvers, were performed in [17]. This variant of the Cahn–Hilliard–Oono equation can also be coupled with the incompressible

---

<sup>3</sup>A better name would be Cahn–Hilliard–Oono–Puri equation; we will however keep the customary one.

Navier–Stokes equations to model a chemically reacting binary fluid (see [246, 247]; see also [48] for the mathematical analysis).

(iii) Proliferation term. In that case,

$$g(x, s) = g(s) = \lambda s(s - 1), \quad \lambda > 0.$$

This function was proposed in [266] in view of biological applications and, more precisely, to model wound healing and tumor growth (in one space dimension; in that case, one can think of a propagation front) and the clustering of malignant brain tumor cells (in two space dimensions); see also [396] for other quadratic functions with chemical applications and [18] for other polynomials with biological applications.

(iv) Fidelity term. In that case,

$$g(x, s) = \lambda_0 \chi_{\Omega \setminus D}(x)(s - h(x)), \quad \lambda_0 > 0, \quad D \subset \Omega, \quad h \in L^2(\Omega),$$

where  $\chi$  denotes the indicator function, and we consider the following equation:

$$\frac{\partial u}{\partial t} + \varepsilon \Delta^2 u - \frac{1}{\varepsilon} \Delta f(u) + g(x, u) = 0, \quad \varepsilon > 0.$$

Written in this way,  $\varepsilon$  corresponds to the interface thickness. This function  $g$  was proposed in [28, 29] in view of applications to binary image inpainting (i.e., black and white images). Here,  $h$  is a given (damaged) image and  $D$  is the inpainting (i.e., damaged) region. Furthermore, the fidelity term  $g(x, u)$  is added in order to keep the solution  $u$  close to the image outside the inpainting region. The idea in this model is to solve the equation up to steady state to obtain an inpainted (i.e., restored) version  $u(x)$  of  $h(x)$ .

The generalized equation (1.17) was studied in [318, 322] (see also [157]) under very general assumptions on the additional term  $g$ , when endowed with Dirichlet boundary conditions. In that case, one essentially recovers the results (well-posedness, regularity and existence of finite-dimensional attractors) known for the original Cahn–Hilliard equation. The case of Neumann boundary conditions is much more involved, due to the fact that one no longer has the conservation of mass, i.e., of the spatial average of the order parameter, when compared with the original Cahn–Hilliard equation with Neumann boundary conditions (see [79, 80, 90, 157, 158, 197]).

Another variant of the Cahn–Hilliard equation is concerned with higher-order Cahn–Hilliard models. More precisely, G. Caginalp and E. Esenturk recently proposed in [60] (see also [76]) higher-order phase-field models in order to account for anisotropic interfaces (see also [274, 387, 407] for other approaches which, however, do not provide an explicit way to compute the anisotropy). More precisely, these authors proposed the following modified free energy, in which we omit the temperature:

$$(1.21) \quad \Psi_{\text{HOGL}} = \int_{\Omega} \left( \frac{1}{2} \sum_{i=1}^M \sum_{|k|=i} a_k |\mathcal{D}^k u|^2 + F(u) \right) dx, \quad M \in \mathbb{N},$$

where, for  $k = (k_1, k_2, k_3) \in (\mathbb{N} \cup \{0\})^3$ ,

$$|k| = k_1 + k_2 + k_3$$

and, for  $k \neq (0, 0, 0)$ ,

$$\mathcal{D}^k = \frac{\partial^{|k|}}{\partial x_1^{k_1} \partial x_2^{k_2} \partial x_3^{k_3}}$$

(we agree that  $\mathcal{D}^{(0,0,0)}v = v$ ). The corresponding higher-order Cahn–Hilliard equation then reads

$$(1.22) \quad \frac{\partial u}{\partial t} - \Delta \sum_{i=1}^M (-1)^i \sum_{|k|=i} a_k \mathcal{D}^{2k} u - \Delta f(u) = 0.$$

For  $M = 1$  (anisotropic Cahn–Hilliard equation), we have an equation of the form

$$\frac{\partial u}{\partial t} + \Delta \sum_{i=1}^3 a_i \frac{\partial^2 u}{\partial x_i^2} - \Delta f(u) = 0$$

and, for  $M = 2$  (sixth-order anisotropic Cahn–Hilliard equation), we have an equation of the form

$$\frac{\partial u}{\partial t} - \Delta \sum_{i,j=1}^3 a_{ij} \frac{\partial^4 u}{\partial x_i^2 \partial x_j^2} + \Delta \sum_{i=1}^3 b_i \frac{\partial^2 u}{\partial x_i^2} - \Delta f(u) = 0.$$

We studied in [86, 312] the corresponding higher-order isotropic model, namely,

$$(1.23) \quad \frac{\partial u}{\partial t} - \Delta P(-\Delta)u - \Delta f(u) = 0,$$

where

$$P(s) = \sum_{i=1}^M a_i s^i, \quad a_k > 0, \quad M \geq 1, \quad s \in \mathbb{R},$$

and, in [87], the anisotropic higher-order model (1.22) (there, numerical simulations were also performed to illustrate the effects of the higher-order terms and of the

anisotropy). Furthermore, these models contain sixth-order Cahn–Hilliard models. We can note that there is currently a strong interest in the study of sixth-order Cahn–Hilliard equations. Such equations arise in situations such as strong anisotropy effects being taken into account in phase separation processes (see [393]), atomistic models of crystal growth (see [28, 29, 153, 186]), the description of growing crystalline surfaces with small slopes which undergo faceting (see [377]), oil-water-surfactant mixtures (see [215, 216]) and mixtures of polymer molecules (see [121]). We refer the reader to [74, 223, 227, 228, 229, 243, 276, 277, 304, 305, 317, 319, 320, 321, 323, 350, 351, 356, 357, 398, 399, 411] for the mathematical and numerical analysis of such models.

We can also note that the variant (1.17) can be relevant in the context of higher-order models (we can mention, for instance, anisotropic effects in tumor growth). We refer the reader to [88] for the analysis and numerical simulations of such models.

We finally mention several other important generalizations and variants of the Cahn–Hilliard equation.

A first one consists in studying systems of Cahn–Hilliard equations to describe phase separation in multicomponent alloys (see [56, 96, 112, 148, 149, 155, 191, 192, 193, 330]). Note that the Cahn–Hilliard equation can be rewritten, equivalently, as a system of two (Cahn–Hilliard) equations. Let us indeed denote by  $A$  and  $B$  the two components and consider, with obvious notation, the free energy

$$\Psi_{\Omega}(u_A, \nabla u_A, u_B, \nabla u_B) = \frac{1}{2} \int_{\Omega} \left( \frac{\alpha}{2} |\nabla u_A|^2 + \frac{\alpha}{2} |\nabla u_B|^2 + F(u_A) + F(u_B) \right) dx.$$

Then, the Cahn–Hilliard system (1.1) is equivalent, again with obvious notation and noting that  $f$  is an odd function in both cases of interest, to

$$\begin{aligned} \frac{\partial u_A}{\partial t} &= \kappa \Delta \mu_A, \quad \mu_A = \frac{1}{2} (-\alpha \Delta u_A + f(u_A)) (= \partial_{u_A} \Psi_{\Omega}), \\ \frac{\partial u_B}{\partial t} &= \kappa \Delta \mu_B, \quad \mu_B = \frac{1}{2} (-\alpha \Delta u_B + f(u_B)) (= \partial_{u_B} \Psi_{\Omega}), \\ u_A + u_B &= 0, \quad \mu_A + \mu_B = 0. \end{aligned}$$

Furthermore, we can see that

$$\mu_A - \mu_B = -\alpha \Delta u_A + f(u_A).$$

We also mention the stochastic Cahn–Hilliard equation (also called the Cahn–Hilliard–Cook equation) which takes into account thermal fluctuations (see [33, 34, 37, 38, 68, 71, 116, 118, 123, 124, 142, 172, 217, 218, 233, 376]).

Then, an important generalization of the Cahn–Hilliard equation is the viscous Cahn–Hilliard equation,

$$-\epsilon\Delta\frac{\partial u}{\partial t} + \frac{\partial u}{\partial t} + \alpha\kappa\Delta^2u - \kappa\Delta f(u) = 0, \quad \epsilon > 0,$$

proposed by A. Novick–Cohen in [344] to account for viscosity effects in the phase separation of polymer/polymer systems (see also [20, 70, 98, 151]). The viscous Cahn–Hilliard equation can also be seen as a particular case of the generalizations proposed by M. Gurtin in [235] (which, in particular, also account for anisotropy) and which are based on a microforce balance, i.e., a new balance law for interactions at a microscopic level (see [41, 42, 44, 85, 139, 140, 211, 221, 314, 315, 316, 325, 326, 331, 336, 367, 368, 369, 408] for the mathematical analysis); we also refer the reader to yet another approach proposed by P. Podio–Guidugli in [359] and studied in, e.g., [102, 103, 104, 105, 110].

Another important generalization of the Cahn–Hilliard equation is the hyperbolic relaxation of the equation,

$$\epsilon\frac{\partial^2 u}{\partial t^2} + \frac{\partial u}{\partial t} + \alpha\kappa\Delta^2u - \kappa\Delta f(u) = 0, \quad \epsilon > 0,$$

proposed in [187, 188, 189, 190, 281] to model the early stages of spinodal decomposition in certain glasses (see also [43, 200, 201, 224, 225, 226, 375] for the mathematical analysis and [373, 374] for the hyperbolic relaxation of the Cahn–Hilliard–Oono equation in the whole space). Actually, the hyperbolic relaxation of the equation is a particular case of more general memory relaxations (for an exponentially decreasing memory kernel) which were studied, e.g., in [111, 113, 115, 202, 203] (see also [361]).

We also mention the convective Cahn–Hilliard equation,

$$\frac{\partial u}{\partial t} + \alpha\kappa\Delta^2u + u \cdot \nabla u - \kappa\Delta f(u) = 0,$$

which describes the dynamics of driven systems such as faceting of growing thermodynamically unstable crystal surfaces (see [135, 136, 137, 213, 291, 405] for the mathematical analysis).

It is important to note that, in realistic physical systems, quenches are usually carried out over a finite period of time, so that phase separation can begin before the final quenching is reached. It is thus important to consider nonisothermal Cahn–Hilliard models. Such models were derived and studied in [12, 13, 183, 184, 329, 382].

The Cahn–Hilliard equation can be coupled with the Allen–Cahn equation which describes the ordering of atoms during the phase separation process (see [11]). This problem was studied, e.g., in [32, 120, 293, 327, 328, 346, 422].

It can also be coupled with the equations for elasticity or viscoelasticity, to account for mechanical effects (see, e.g., [16, 31, 39, 40, 69, 128, 192, 193, 194, 314, 315, 352, 353, 354, 355, 366]).

We also mention the coupling of the Cahn–Hilliard equation with the Navier–Stokes equations in the context of two-phase (multiphase) flows (see, e.g., [2, 4, 48, 49, 50, 51, 52, 54, 67, 64, 84, 91, 99, 167, 168, 179, 180, 182, 210, 236, 242, 255, 267, 270, 272, 279, 292, 299, 332, 419, 423]) and some related models such as the Cahn–Hilliard–Hele–Shaw and Cahn–Hilliard–Brinkman equations (see, e.g., [47, 114, 125, 126, 127, 162, 209, 240, 402, 404, 409, 418]). Related models can also be used to model tumor growth (see, e.g., [101, 106, 107, 108, 119, 169, 195, 196, 257, 298]).

We finally refer the reader to, e.g., [1, 9, 10, 14, 15, 20, 21, 22, 23, 24, 25, 26, 36, 45, 51, 52, 53, 54, 55, 57, 58, 65, 66, 73, 75, 77, 92, 94, 117, 129, 130, 144, 145, 146, 150, 156, 159, 160, 161, 163, 169, 170, 171, 199, 214, 219, 220, 222, 234, 237, 238, 239, 241, 248, 249, 256, 262, 263, 265, 267, 268, 269, 270, 271, 272, 282, 283, 284, 285, 286, 288, 289, 295, 301, 306, 313, 338, 339, 364, 371, 380, 381, 385, 386, 392, 400, 401, 403, 406, 410, 413, 414, 416, 425, 426] for the numerical analysis and simulations of the Cahn–Hilliard equation (and several of its generalizations). Note that, as suggested in [146], it is in general preferable to build numerical schemes for the Cahn–Hilliard system (1.1) rather than the equivalent fourth-order in space Cahn–Hilliard equation. This has indeed the advantage of splitting the fourth-order equation into a system of two second-order ones which is easier to deal with.

**Notation.** We denote by  $\|\cdot\|$  the usual norms on  $L^2(\Omega)$  and  $L^2(\Omega)^n$  (with associated scalar product  $((\cdot, \cdot))$ ). More generally,  $\|\cdot\|_X$  denotes the norm on the Banach space  $X$ .

In what follows,  $\Omega$  is a bounded and regular (as regular as needed) domain of  $\mathbb{R}^n$ ,  $n = 1, 2$  or  $3$ .

## 2. LINEAR OPERATORS [388]

We consider the spaces  $L^2(\Omega)$  and  $H^1(\Omega)$  which, endowed with their usual scalar products and associated norms, are Hilbert spaces.

Of course,  $(v, v) \mapsto ((\nabla u, \nabla v))$  is not a scalar product in  $H^1(\Omega)$ , as it is not coercive.

To overcome this, we set

$$\langle u \rangle = \frac{1}{\text{Vol}(\Omega)} \int_{\Omega} u \, dx, \quad u \in L^1(\Omega),$$

$$\langle u \rangle = \frac{1}{\text{Vol}(\Omega)} \langle u, 1 \rangle, \quad u \in H^{-1}(\Omega),$$

where  $H^{-1}(\Omega)$  is the topological dual of  $H^1(\Omega)$ ,  $H^{-1}(\Omega) = H^1(\Omega)'$ , and  $\langle \cdot, \cdot \rangle$  denotes the duality pairing between  $H^{-1}(\Omega)$  and  $H^1(\Omega)$ .

We then set

$$H = \dot{L}^2(\Omega) = \{u \in L^2(\Omega), \langle u \rangle = 0\},$$

$$V = \dot{H}^1(\Omega) = H^1(\Omega) \cap H.$$

These spaces are also Hilbert spaces, when endowed with the induced scalar products. Furthermore,  $((\cdot, \cdot))_V = ((\nabla \cdot, \nabla \cdot))$  is a scalar product on  $V$ , with associated norm  $\|\cdot\|_V$ , which is equivalent to the usual  $H^1$ -scalar product (owing to the Poincaré-Wirtinger inequality; see below).

Let  $V' = \{u \in H^{-1}(\Omega), \langle u \rangle = 0\}$  be the topological dual of  $V$ . Then, we know from Riesz' representation theorem that,  $\forall l \in V'$ , there exists a unique  $u \in V$  such that  $((u, v))_V = \langle l, v \rangle$ ,  $\forall v \in V$ , where  $\langle \cdot, \cdot \rangle$  also denotes the duality pairing between  $V'$  and  $V$ .

Identifying  $H$  with its topological dual  $H'$ , we have the Hilbert triplet  $V \subset H \equiv H' \subset V'$ , with dense, continuous and compact injections. Furthermore, if  $u$  and  $v$  are in  $H$  and  $V$ , respectively, then  $\langle u, v \rangle = ((u, v))_H$ .

We can then define the linear operator  $A : V \rightarrow V'$  by

$$\langle Au, v \rangle = ((u, v))_V, \quad \forall u, v \in V.$$

This operator is an isomorphism from  $V$  onto  $V'$ .

We now set

$$D(A) = A^{-1}(H) = \{u \in V, Au \in H\} = \{u \in \dot{H}^1(\Omega), -\Delta u \in \dot{L}^2(\Omega)\},$$

called domain of  $A$ . Note indeed that, if  $((u, v))_V = ((f, v))$ ,  $\forall v \in V$  and for  $f \in H$ , then

$$((\nabla u, \nabla v)) = ((f, v)), \quad \forall v \in H^1(\Omega)$$

(it suffices to replace  $v$  by  $v - \langle v \rangle$ ). Taking then  $v \in \mathcal{D}(\Omega) \equiv \mathcal{C}_c^\infty(\Omega)$ , it is easy to see that  $-\Delta u = f$  in the sense of distributions and thus in  $L^2(\Omega)$ .

Next, since  $u \in H^1(\Omega)$  and  $\Delta u \in L^2(\Omega)$ , the trace  $\frac{\partial u}{\partial \nu}$  can be defined in  $H^{-\frac{1}{2}}(\Gamma)$  and a generalized form of Green's formula is valid for every  $v \in H^1(\Omega)$  (see [389]; see also [388], Chapter II, Example 2.5), yielding

$$-((\Delta u, v)) = -\left\langle \frac{\partial u}{\partial \nu}, v \right\rangle_{H^{-\frac{1}{2}}(\Gamma), H^{\frac{1}{2}}(\Gamma)} + ((\nabla u, \nabla v)), \quad \forall v \in H^1(\Omega).$$

We thus deduce that

$$\left\langle \frac{\partial u}{\partial \nu}, v \right\rangle_{H^{-\frac{1}{2}}(\Gamma), H^{\frac{1}{2}}(\Gamma)} = 0, \quad \forall v \in H^1(\Omega),$$

so that

$$\frac{\partial u}{\partial \nu} = 0, \text{ on } \Gamma$$

(in  $H^{-\frac{1}{2}}(\Gamma)$ ). Having this, it follows from classical elliptic regularity results (see [6, 7, 8]) that  $u \in H^2(\Omega)$ . Finally,

$$D(A) = \{u \in H^2(\Omega) \cap V, \frac{\partial u}{\partial \nu} = 0 \text{ on } \Gamma\}$$

and  $Au = f$ ,  $u \in D(A)$  and  $f \in H$ , is equivalent to

$$-\Delta u = f, \text{ in } \Omega, \frac{\partial u}{\partial \nu} = 0, \text{ on } \Gamma.$$

**Remark 2.1.** More generally, if  $f \in H^m(\Omega)$ ,  $m \geq 0$ , then  $u \in H^{m+2}(\Omega)$ .

**2.1. Spectral properties of the operator  $A$ .** First note that  $A$  is selfadjoint (since  $((\cdot, \cdot))_V$  is symmetric). Furthermore, since  $V \subset H$  is compact, then  $A^{-1} : H \rightarrow H$  is compact (and selfadjoint). Indeed,  $A^{-1} : H \rightarrow D(A)$  is continuous, so that  $A^{-1} : H \rightarrow V$  is also continuous (note that it follows from the regularity mentioned above that the injection  $D(A) \subset V$  is continuous) and we conclude owing to the compact injection  $V \subset H$ .

We thus conclude that  $A^{-1}$  is compact, selfadjoint and positive (as an operator in  $H$ ). Therefore, there exists an orthonormal basis  $(w_j)$ ,  $j \in \mathbb{N}$ , of  $H$  formed of eigenvectors of  $A$ :

$$A^{-1}w_j = \mu_j w_j, \mu_j \rightarrow 0 \text{ as } j \rightarrow +\infty, \mu_j > 0.$$

Since  $w_j = \frac{1}{\mu_j} A^{-1}w_j \in D(A)$ , then

$$Aw_j = \lambda_j w_j, \lambda_j = \frac{1}{\mu_j},$$

and  $w_j$ ,  $\lambda_j$  are eigenvectors/eigenvalues of  $A$ , where  $0 < \lambda_1 \leq \lambda_2 \leq \dots$ ,  $\lambda_j \rightarrow +\infty$  as  $j \rightarrow +\infty$  (note that  $\langle Aw_j, w_j \rangle = \lambda_j \|w_j\|^2 > 0$ ). Furthermore, the  $w_j$ 's are orthogonal in  $V$  for  $((\cdot, \cdot))_V$ . Indeed, if  $j \neq k$ , then

$$((w_j, w_k))_V = \langle Aw_j, w_k \rangle = \lambda_j ((w_j, w_k)) = 0.$$

However, this family is not orthonormal, since

$$\langle Aw_j, w_j \rangle = ((w_j, w_j))_V = \lambda_j \|w_j\|^2 = \lambda_j.$$

We can now compute the norms of  $u$  in terms of the spectral elements of  $A$ . Indeed:

- If  $u \in H$ ,  $u = \sum_{j=1}^{\infty} u_j w_j$ , then  $\|u\|^2 = \sum_{j=1}^{\infty} |u_j|^2$ .



• If  $u \in V$ , then  $\|u\|_V^2 = ((\sum_{j=1}^{\infty} u_j w_j, \sum_{j=1}^{\infty} u_j w_j))_V = \sum_{j=1}^{\infty} |u_j|^2 ((w_j, w_j))_V = \sum_{j=1}^{\infty} \lambda_j |u_j|^2$ . We also have

$$V = \{u \in H, u = \sum_{j=1}^{\infty} u_j w_j, \sum_{j=1}^{\infty} \lambda_j |u_j|^2 < +\infty\}.$$

• If  $u \in D(A)$ , then  $Au = A(\sum_{j=1}^{\infty} u_j w_j) = \sum_{j=1}^{\infty} u_j A w_j = \sum_{j=1}^{\infty} \lambda_j u_j w_j$  (the sequence converges) and  $\|u\|_{D(A)}^2 = \|Au\|^2 = \sum_{j=1}^{\infty} \lambda_j^2 |u_j|^2$  (graph norm). We also have

$$D(A) = \{u \in H, u = \sum_{j=1}^{\infty} u_j w_j, \sum_{j=1}^{\infty} \lambda_j^2 |u_j|^2 < +\infty\}.$$

Furthermore, on  $D(A)$ , the graph norm  $\|A \cdot\|$  is equivalent to the usual  $H^2$ -norm, owing to the elliptic regularity results mentioned above.

**2.2. The spaces  $D(A^\alpha)$  and the operators  $A^\alpha$ .** We first consider the case  $\alpha \geq 0$ .

By convention,  $D(A^0) = H$  and  $A^0 = I$  (identity operator). Then, for  $\alpha > 0$ , we set (see also above)

$$D(A^\alpha) = \{u \in H, u = \sum_{j=1}^{\infty} u_j w_j, \sum_{j=1}^{\infty} \lambda_j^{2\alpha} |u_j|^2 < +\infty\}$$

which we endow with the graph norm

$$\|u\|_{D(A^\alpha)} = \left( \sum_{j=1}^{\infty} \lambda_j^{2\alpha} |u_j|^2 \right)^{\frac{1}{2}}$$

and the associated scalar product

$$((u, v))_{D(A^\alpha)} = \sum_{j=1}^{\infty} \lambda_j^{2\alpha} u_j v_j, \quad u = \sum_{j=1}^{\infty} u_j w_j, \quad v = \sum_{j=1}^{\infty} v_j w_j.$$

Endowed with this scalar product,  $D(A^\alpha)$  is a Hilbert space.

The operator  $A^\alpha : D(A^\alpha) \rightarrow H$  is then defined by

$$A^\alpha u = \sum_{j=1}^{\infty} \lambda_j^\alpha u_j w_j, \quad u = \sum_{j=1}^{\infty} u_j w_j.$$

**Example 2.2.** (i) We recover the above convention and definition for  $\alpha = 0$  and  $\alpha = 1$ .

(ii) For  $\alpha = \frac{1}{2}$ ,  $D(A^{\frac{1}{2}}) = V$ .

(iii) For  $\alpha = 2$ ,

$$D(A^2) = \{u \in H, u = \sum_{j=1}^{\infty} u_j w_j, \sum_{j=1}^{\infty} \lambda_j^4 |u_j|^2 < +\infty\} = \{u \in D(A), Au \in D(A)\}.$$

It thus follows for the elliptic regularity results of [6, 7, 8] that

$$D(A^2) = \{u \in H^4(\Omega) \cap V, \frac{\partial u}{\partial \nu} = \frac{\partial \Delta u}{\partial \nu} = 0 \text{ on } \Gamma\}.$$

Note that we recover the two (Neumann) boundary conditions associated with the Cahn–Hilliard equation. Indeed,  $Au = f$ ,  $u \in D(A^2)$  and  $f \in H$  is equivalent to

$$\Delta^2 u = f, \text{ in } \Omega, \frac{\partial u}{\partial \nu} = \frac{\partial \Delta u}{\partial \nu} = 0, \text{ on } \Gamma.$$

Furthermore, the graph norm  $\|A^2 \cdot\|$  is equivalent to the usual  $H^4$ -norm on  $D(A^2)$ .

**Remark 2.3.** If  $\alpha > \alpha' \geq 0$ , then  $D(A^\alpha) \subset D(A^{\alpha'})$  with continuous, dense and compact injection.

We now turn to the case  $\alpha < 0$ .

We consider, on  $H$ , the continuous norm  $(\sum_{j=1}^{\infty} \lambda_j^{2\alpha} |u_j|^2)^{\frac{1}{2}}$ ,  $u = \sum_{j=1}^{\infty} u_j w_j$ . Then, by definition,  $D(A^\alpha)$  is the closure of  $H$  for this norm.

We thus have  $H \subset D(A^\alpha)$  with continuous injection (this injection is indeed continuous, since  $\sum_{j=1}^{\infty} \lambda_j^{2\alpha} |u_j|^2 \leq \lambda_1^{2\alpha} \sum_{j=1}^{\infty} |u_j|^2$ ).

**Remark 2.4.** a) If  $\alpha > \alpha'$ ,  $\alpha, \alpha' \in \mathbb{R}$ , then  $D(A^\alpha) \subset D(A^{\alpha'})$  with continuous, dense and compact injection. Furthermore, if  $\alpha > 0$ , then  $D(A^\alpha)' = D(A^{-\alpha})$  (topological dual).

b) We could have also defined  $D(A^\alpha)$ ,  $\alpha < 0$ , as the topological dual of  $D(A^{-\alpha})$ .

The operator  $A^\alpha$ ,  $\alpha < 0$ , is then defined as above,  $A^\alpha : D(A^\alpha) \rightarrow H$ .

**Example 2.5.** (i) For  $\alpha = -1$ ,  $D(A^{-1}) = D(A)'$ , as mentioned above.

(ii) For  $\alpha = -\frac{1}{2}$ ,  $D(A^{-\frac{1}{2}}) = D(A^{\frac{1}{2}})' = V'$  and the graph norm,  $\|A^{-\frac{1}{2}} \cdot\|$ , is equivalent to the usual  $H^{-1}$ -norm on  $D(A^{-\frac{1}{2}})$ .

**Remark 2.6.** The norm

$$u \mapsto (\|A^{-\frac{1}{2}}(u - \langle u \rangle)\|^2 + \langle u \rangle^2)^{\frac{1}{2}}$$

is a norm on  $H^{-1}(\Omega)$  which is equivalent to the usual  $H^{-1}$ -norm.

We can note that  $D(A) \subset D(A^{\frac{1}{2}}) \equiv V \subset D(A^0) \equiv H \subset D(A^{-\frac{1}{2}}) \equiv V'$ , with continuous, dense and compact injections. Furthermore,

$$\|u\|_V^2 = \sum_{j=1}^{+\infty} \lambda_j |u_j|^2 \geq \lambda_1 \sum_{j=1}^{+\infty} |u_j|^2 = \lambda_1 \|u\|^2,$$

which yields the Poincaré type inequality

$$\|u\| \leq \frac{1}{\sqrt{\lambda_1}} \|u\|_V, \quad \forall u \in V.$$

Note that we can obtain similar estimates on the norms of all injections.

**Remark 2.7.** We deduce from the above the Poincaré–Wirtinger inequality

$$(2.1) \quad \|u - \langle u \rangle\| \leq \frac{1}{\sqrt{\lambda_1}} \|u\|_V, \quad \forall u \in H^1(\Omega).$$

**2.3. The linear equation**  $\frac{du}{dt} + A^2u = f(t)$ . The linear equation associated with the Cahn–Hilliard equation reads

$$\begin{aligned} \frac{\partial u}{\partial t} + \Delta^2 u &= f(x, t), \\ \frac{\partial u}{\partial \nu} = \frac{\partial \Delta u}{\partial \nu} &= 0 \text{ on } \Gamma. \end{aligned}$$

Having in mind the operator  $A$  defined in the previous subsection, we can rewrite this boundary value problem in the following functional form:

$$\frac{du}{dt} + A^2u = f(t).$$

In particular, if  $f \in L^2(0, T; H)$ ,  $u \in L^2(0, T; D(A^2))$  and  $\frac{du}{dt} \in L^2(0, T; H)$ ,  $T > 0$  given, this equation makes sense in  $L^2(0, T; H)$ . Furthermore, it is associated with the variational formulation

Find  $u : [0, T] \rightarrow D(A)$  such that

$$\frac{d}{dt}((u, v)) + ((Au, Av)) = ((f(t), v)), \quad \forall v \in D(A),$$

in the sense of distributions.

We have the following result (existence and uniqueness of weak solutions).

**Theorem 2.8.** *We assume that  $f \in L^2(0, T; D(A^{-1}))$  and  $u_0 \in H$ ,  $T > 0$  given. Then, the linear initial value problem*

$$(2.2) \quad \frac{du}{dt} + A^2u = f(t) \text{ in } \mathcal{D}'(0, T; D(A^{-1})),$$

$$(2.3) \quad u(0) = u_0 \text{ in } H,$$

possesses a unique solution  $u$  such that  $u \in \mathcal{C}([0, T]; H) \cap L^2(0, T; D(A))$  and  $\frac{du}{dt} \in L^2(0, T; D(A^{-1}))$ .

**Remark 2.9.** Note that, since  $D(A) \subset H \subset D(A^{-1}) \equiv D(A)'$  with continuous injections, then it follows from Lions' theorem (see [290]; see also the next subsection below) that, if  $u \in L^2(0, T; D(A))$  and  $\frac{du}{dt} \in L^2(0, T; D(A^{-1}))$ , then  $u \in \mathcal{C}([0, T]; H)$ . Therefore,  $u(0)$  makes sense in  $H$ .

*Proof. Uniqueness:*

Let  $u_1$  and  $u_2$  be solutions to

$$\begin{aligned} \frac{du_1}{dt} + A^2u_1 &= f_1(t) \text{ in } \mathcal{D}'(0, T; D(A^{-1})), \\ u_1(0) &= u_{0,1} \text{ in } H, \end{aligned}$$

and

$$\begin{aligned} \frac{du_2}{dt} + A^2u_2 &= f_2(t) \text{ in } \mathcal{D}'(0, T; D(A^{-1})), \\ u_2(0) &= u_{0,2} \text{ in } H. \end{aligned}$$

Then,  $u = u_1 - u_2$ ,  $f = f_1 - f_2$  and  $u_0 = u_{1,0} - u_{2,0}$  satisfy

$$(2.4) \quad \frac{du}{dt} + A^2u = f(t) \text{ in } \mathcal{D}'(0, T; D(A^{-1})),$$

$$(2.5) \quad u(0) = u_0 \text{ in } H.$$

The above equation is associated with the variational formulation

$$(2.6) \quad \frac{d}{dt}((u, v)) + ((Au, Av)) = \langle f(t), v \rangle \text{ in } \mathcal{D}'(0, T), \quad \forall v \in D(A),$$

where  $\langle \cdot, \cdot \rangle$  denotes the duality product in  $D(A^{-1})$ .

Taking  $v = u$  in (2.6), we have

$$\frac{1}{2} \frac{d}{dt} \|u\|^2 + \|Au\|^2 = \langle f(t), u \rangle,$$

which yields, noting that

$$|\langle f(t), u \rangle| \leq \|f(t)\|_{D(A^{-1})} \|Au\| \leq \frac{1}{2} \|f(t)\|_{D(A^{-1})}^2 + \frac{1}{2} \|Au\|^2$$

the differential inequality

$$\frac{d}{dt} \|u\|^2 + \|Au\|^2 \leq \|f(t)\|_{D(A^{-1})}^2,$$

so that, integrating over  $[0, t]$ ,  $0 \leq t \leq T$ ,

$$\|u(t)\|^2 \leq \|u_0\|^2 + \|f\|_{L^2(0, T; D(A^{-1}))}^2.$$

This gives the continuous dependence with respect to the data (i.e.,  $(u_0, f) \mapsto u(t)$  is continuous from  $H \times L^2(0, T; D(A^{-1}))$  onto  $H$ ), as well as the uniqueness (taking  $f_1 = f_2$  and  $u_{1,0} = u_{2,0}$ ).

**Existence:**

The proof of existence is based on a standard Galerkin scheme.

We consider, for  $m \in \mathbb{N}$  given, the following approximated problem:

Find  $u_m = \sum_{i=1}^m u_i w_i$ ,  $u_i = u_i(t)$ ,  $i = 1, \dots, m$ , such that

$$(2.7) \quad \frac{d}{dt} ((u_m, v)) + ((Au_m, Av)) = \langle f(t), v \rangle, \quad \forall v \in W_m \equiv \text{Span}(w_1, \dots, w_m),$$

$$(2.8) \quad u_m(0) = u_{0,m},$$

where  $u_{0,m}$  is the projection (in  $H$ ) of  $u_0$  onto  $W_m$ ,

$$u_{0,m} = \sum_{i=1}^m ((u_0, w_i)) w_i.$$

Here, the  $w_j$ 's are eigenfunctions of  $A$  as constructed above. Note that the  $u_i$ 's do not depend on  $m$ , since the problem is linear. Furthermore, (2.7) is equivalent to (taking  $v = w_i$ ,  $i = 1, \dots, m$ , in (2.7))

$$\frac{du_i}{dt} + \lambda_i^2 u_i = \langle f(t), w_i \rangle, \quad i = 1, \dots, m,$$

so that

$$u_i(t) = e^{-\lambda_i^2 t} u_{i,0} + e^{-\lambda_i^2 t} \int_0^t e^{\lambda_i^2 s} \langle f(s), w_i \rangle ds, \quad i = 1, \dots, m,$$

where  $u_{i,0} = ((u_0, w_i))$ .

We thus have the existence of the maximal solution  $u_m$  to (2.7); actually, this solution is global and defined on  $[0, T]$ .

We now take  $v = w_i$  in (2.7) and multiply the resulting equation by  $u_i$  to find

$$u_i \frac{du_i}{dt} + \lambda_i^2 |u_i| = \langle f(t), u_i w_i \rangle.$$

Summing over  $i = 1, \dots, m$  and noting that  $u_i \frac{du_i}{dt} = \frac{1}{2} \frac{d}{dt} |u_i|^2$ , we obtain

$$\frac{1}{2} \frac{d}{dt} \|u_m\|^2 + \|Au_m\|^2 = \langle f(t), u_m \rangle,$$

which yields, proceeding as above,

$$\frac{d}{dt} \|u_m\|^2 + \|Au_m\|^2 \leq \|f(t)\|_{D(A^{-1})}^2.$$

Therefore,  $u_m$  is bounded, independently of  $m$ , in  $L^2(0, T; D(A))$  and  $L^\infty(0, T; H)$ . Hence, up to a subsequence which we do not relabel, there exists  $u \in L^\infty(0, T; H) \cap L^2(0, T; D(A))$  such that  $u_m \rightarrow u$  in  $L^\infty(0, T; H)$  weak star and in  $L^2(0, T; D(A))$  weak. This allows to pass to the limit in (2.7) and finish the proof of existence (the regularity on  $\frac{du}{dt}$  immediately follows from the equation and the regularity on  $u$  and  $f$ ). □

We also have the following result (existence and uniqueness of strong solutions).

**Theorem 2.10.** *We assume that  $u_0 \in D(A)$  and  $f \in L^2(0, T; H)$ . Then, the solution  $u$  to (2.2) given in Theorem 2.8 satisfies  $u \in \mathcal{C}([0, T]; D(A)) \cap L^2(0, T; D(A^2))$  and  $\frac{du}{dt} \in L^2(0, T; H)$  and is a strong solution to (2.2) (i.e., (2.2) holds in  $L^2(0, T; H)$ ).*

**Remark 2.11.** Note that  $D(A^2) \subset D(A) \subset H$  with continuous injections. Therefore,  $u \in L^2(0, T; D(A^2))$  and  $\frac{du}{dt} \in L^2(0, T; H)$  imply that  $u \in \mathcal{C}([0, T]; D(A))$ . Therefore,  $u(0)$  makes sense in  $D(A)$ .

*Proof.* Owing to the regularity on  $f$ , (2.6) and (2.7) now read

$$(2.9) \quad \frac{d}{dt}((u, v)) + ((Au, Av)) = ((f(t), v)) \text{ in } \mathcal{D}'(0, T), \quad \forall v \in D(A),$$

and

$$(2.10) \quad \frac{d}{dt}((u_m, v)) + ((Au_m, Av)) = ((f(t), v)), \quad \forall v \in W_m,$$

respectively.

We take  $v = w_i$  in (2.10) and multiply the resulting equation by  $\lambda_i^2 u_i$ . This yields

$$\lambda_i^2 u_i \frac{du_i}{dt} + \lambda_i^4 |u_i| = ((f(t), \lambda_i^2 u_i w_i)).$$

Summing over  $i = 1, \dots, m$ , we have

$$\frac{1}{2} \frac{d}{dt} \|Au_m\|^2 + \|A^2u_m\|^2 = ((f(t), Au_m)),$$

Noting that

$$|((f(t), Au_m))| \leq \|f(t)\| \|Au_m\| \leq \frac{1}{c_0} \|f(t)\| \|A^2u_m\|,$$

where  $c_0$  is such that  $\|A^2u_m\| \geq c_0 \|Au_m\|$  and depends on  $\lambda_1$  (see the previous subsection), this yields

$$\frac{d}{dt} \|Au_m\|^2 + \|A^2u_m\|^2 \leq \frac{1}{c_0^2} \|f(t)\|^2.$$

Therefore,  $u_m$  is bounded, independently of  $m$ , in  $L^2(0, T; D(A))$  and  $L^2(0, T; D(A^2))$  and we can pass to the limit as above.  $\square$

Note that, if  $u \in L^\infty(0, T; H) \cap L^2(0, T; D(A))$  is solution to (2.2), then

$$A^{-1} \frac{du}{dt} + Au = A^{-1}f(t) \text{ in } L^2(0, T; H)$$

and conversely.

We will actually consider a weaker formulation of (2.2), namely,

$$(2.11) \quad A^{-1} \frac{du}{dt} + Au = A^{-1}f(t) \text{ in } L^2(0, T; V'),$$

$$(2.12) \quad u(0) = u_0 \text{ in } V',$$

which is associated with the variational formulation

Find  $u : [0, T] \rightarrow V$  such that

$$(2.13) \quad \frac{d}{dt} ((A^{-1}u, v)) + \langle Au, v \rangle = \langle A^{-1}f(t), v \rangle \text{ in } \mathcal{D}'(0, T), \quad \forall v \in V,$$

where  $\langle \cdot, \cdot \rangle$  denotes the duality product in  $V'$ . Note that  $A^{-1} \frac{du}{dt} = \frac{dA^{-1}u}{dt}$  and recall that  $V' = D(A^{-\frac{1}{2}})$ .

We have the following result (existence and uniqueness of very weak solutions).

**Theorem 2.12.** *We assume that  $u_0 \in V'$  and  $f \in L^2(0, T; D(A^{-\frac{3}{2}}))$ . Then, (2.11)-(2.12) possesses a unique solution  $u$  such that  $u \in \mathcal{C}([0, T]; V') \cap L^2(0, T; V)$  and  $\frac{du}{dt} \in L^2(0, T; D(A^{-\frac{3}{2}}))$ .*

**Remark 2.13.** Note that  $V \subset V' \subset D(A^{-\frac{3}{2}})$  with continuous injections. Therefore,  $u \in L^2(0, T; V)$  and  $\frac{du}{dt} \in L^2(0, T; D(A^{-\frac{3}{2}}))$  imply that  $u \in \mathcal{C}([0, T]; V')$ . Therefore,  $u(0)$  makes sense in  $V'$ . Also note that, if  $f \in L^2(0, T; D(A^{-\frac{3}{2}}))$ , then  $A^{-1}f \in L^2(0, T; V')$ ; the same holds for  $\frac{du}{dt}$ .

*Proof. Uniqueness:*

Let  $u_1$  and  $u_2$  be solutions to

$$\begin{aligned} A^{-1} \frac{du_1}{dt} + Au_1 &= A^{-1} f_1(t) \text{ in } \mathcal{D}'(0, T; V'), \\ u_1(0) &= u_{0,1} \text{ in } V', \end{aligned}$$

and

$$\begin{aligned} A^{-1} \frac{du_2}{dt} + Au_2 &= A^{-1} f_2(t) \text{ in } \mathcal{D}'(0, T; V'), \\ u_2(0) &= u_{0,2} \text{ in } V'. \end{aligned}$$

Then,  $u = u_1 - u_2$ ,  $f = f_1 - f_2$  and  $u_0 = u_{1,0} - u_{2,0}$  satisfy

$$(2.14) \quad A^{-1} \frac{du}{dt} + Au = A^{-1} f(t) \text{ in } \mathcal{D}'(0, T; V'),$$

$$(2.15) \quad u(0) = u_0 \text{ in } V'.$$

The above equation is associated with the variational formulation

$$(2.16) \quad \frac{d}{dt} \langle A^{-1}u, v \rangle + \langle Au, v \rangle = \langle A^{-1}f(t), v \rangle \text{ in } \mathcal{D}'(0, T), \quad \forall v \in V.$$

Taking  $v = u$  in (2.16), we have

$$\frac{1}{2} \frac{d}{dt} \|u\|_{-1}^2 + \|u\|_V^2 = \langle A^{-1}f(t), u \rangle,$$

where  $\|\cdot\|_{-1} = \|A^{-\frac{1}{2}} \cdot\|$ . Noting that  $|\langle A^{-1}f(t), u \rangle| \leq \|f(t)\|_{D(A^{-\frac{3}{2}})} \|u\|_V$ , this yields the differential inequality

$$\frac{d}{dt} \|u\|_{-1}^2 + \|u\|_V^2 \leq \|f(t)\|_{D(A^{-\frac{3}{2}})}^2,$$

so that, integrating over  $[0, t]$ ,  $0 \leq t \leq T$ ,

$$\|u(t)\|_{-1}^2 \leq \|u_0\|_{-1}^2 + \|f\|_{L^2(0, T; D(A^{-\frac{3}{2}}))}^2.$$



This gives the continuous dependence with respect to the data (i.e.,  $(u_0, f) \mapsto u(t)$  is continuous from  $V' \times L^2(0, T; D(A^{-\frac{3}{2}}))$  onto  $V'$ ), as well as the uniqueness.

**Existence:**

We again employ a Galerkin scheme.

We consider, for  $m \in \mathbb{N}$  given, the approximated problem

Find  $u_m = \sum_{i=1}^m u_i w_i$ ,  $u_i = u_i(t)$ ,  $i = 1, \dots, m$ , such that

$$\frac{d}{dt}((A^{-1}u_m, v)) + ((Au_m, v)) = \langle f(t), v \rangle, \quad \forall v \in W_m,$$

$$u_m(0) = u_{0,m},$$

where  $u_{0,m}$  is the projection (in  $V'$ ) of  $u_0$  onto  $W_m$ . Note that here we have enough regularity to write a scalar product in  $H$ , instead of a duality product.

Taking  $v = w_i$ , multiplying the resulting equation by  $u_i$  and summing over  $i = 1, \dots, m$ , we find

$$\frac{1}{2} \frac{d}{dt} \|u_m\|_{-1}^2 + \|u_m\|_V^2 = \langle f(t), u_m \rangle,$$

which yields

$$\frac{d}{dt} \|u_m\|_{-1}^2 + \|u_m\|_V^2 \leq \|f(t)\|_{D(A^{-\frac{3}{2}})}^2.$$

Therefore,  $u_m$  is bounded independently of  $m$  in  $L^\infty(0, T; V')$  and  $L^2(0, T; V)$ . We can then pass to the limit in the approximated problems and finish the proof of the theorem. □

**Remark 2.14.** Assuming more regularity on  $u_0$  and  $f$ , we can obtain more regularity on the solutions and recover the strong formulation, in particular.

### 3. AUBIN–LIONS COMPACTNESS RESULTS [383]

Let  $X_0$ ,  $X$  and  $X_1$  be three Banach spaces such that  $X \subset X_0 \subset X_1$  with dense and continuous injections, where  $X \subset X_0$  is compact. Let  $p \geq 1$ ,  $1 \leq q \leq +\infty$  and  $T > 0$  be given. We set

$$W = \left\{ u \in L^p(0, T; X), \frac{du}{dt} \in L^q(0, T; X_1) \right\}.$$

- If  $p < +\infty$ , then the embedding  $W \in L^p(0, T; X_0)$  is compact.
- If  $p = +\infty$ , then the embedding  $W \in \mathcal{C}([0, T]; X_0)$  is compact.

**Remark 3.1.** When  $p = q = 2$  and the injections are only dense and continuous, we recover Lions' theorem, namely,  $W \subset \mathcal{C}([0, T]; X_0)$  with continuous injection (see [290]; see also [384] for insightful discussions on Lions' theorem).

#### 4. SOME USEFUL INEQUALITIES

**4.1. Interpolation inequalities [388].** Let  $m \in \mathbb{N} \cup \{0\}$  and  $s \in (0, 1)$ . Then,

$$(4.1) \quad \|u\|_{H^{m+s}(\Omega)} \leq c \|u\|_{H^m(\Omega)}^\theta \|u\|_{H^{m+1}(\Omega)}^{1-\theta}, \quad \forall u \in H^{m+1}(\Omega),$$

where  $\theta = 1 - s$  and  $1 - \theta = s$ . It is understood here that  $H^0(\Omega) = L^2(\Omega)$ .

**Example 4.1.** There holds,  $\forall u \in H^1(\Omega)$ ,

$$(4.2) \quad \|u\|_{H^{\frac{1}{2}}(\Omega)} \leq c \|u\|^{\frac{1}{2}} \|u\|_{H^1(\Omega)}^{\frac{1}{2}}.$$

Let  $m_2 > m_1 \geq 0$ ,  $m_1, m_2 \in \mathbb{N} \cup \{0\}$ , and  $s \in (m_1, m_2)$ . Then,

$$(4.3) \quad \|u\|_{H^s(\Omega)} \leq c \|u\|_{H^{m_1}(\Omega)}^\theta \|u\|_{H^{m_2}(\Omega)}^{1-\theta}, \quad \forall u \in H^{m_2}(\Omega),$$

where  $\theta = \frac{m_2 - s}{m_2 - m_1}$  and  $1 - \theta = \frac{s - m_1}{m_2 - m_1}$ .

**Example 4.2.** (i) There holds,  $\forall u \in H^4(\Omega)$ ,

$$(4.4) \quad \|u\|_{H^2(\Omega)} \leq c \|u\|_{H^1(\Omega)}^{\frac{2}{3}} \|u\|_{H^4(\Omega)}^{\frac{1}{3}}.$$

(ii) There holds,  $\forall u \in H^4(\Omega)$ ,

$$(4.5) \quad \|u\|_{H^{\frac{7}{4}}(\Omega)} \leq c \|u\|_{H^1(\Omega)}^{\frac{3}{4}} \|u\|_{H^4(\Omega)}^{\frac{1}{4}}.$$

**Remark 4.3.** a) One important question is to find in an easy way the coefficients in the above interpolation inequalities. This can be done in several ways. One possibility is via a dimensional analysis, based on lengths. Let us consider the interpolation inequality (4.3). In  $\mathbb{R}^n$ , a volume corresponds to the dimension  $L^n$ . Then, a derivative corresponds to  $L^{-1}$ . Thus, for instance, the norm of  $u$  in  $H^{m_1}(\Omega)$  corresponds to  $(L^n L^{-2m_1})^{\frac{1}{2}}$  and, by analogy, the norm of  $u$  in  $H^s(\Omega)$ , even for  $s$  noninteger, to  $(L^n L^{-2s})^{\frac{1}{2}}$ . Therefore, we have, in (4.3),  $(L^n L^{-2s})^{\frac{1}{2}} = (L^n L^{-2m_1})^{\frac{\theta}{2}} (L^n L^{-2m_2})^{\frac{1-\theta}{2}}$ , so that  $s = \theta m_1 + (1 - \theta) m_2$  and the result follows.

b) In particular, we can see that the coefficients are independent of the space dimension  $n$ .

4.2. **Agmon's inequalities** [388]. There holds, in two space dimensions,

$$(4.6) \quad \|u\|_{L^\infty(\Omega)} \leq c \|u\|^{\frac{1}{2}} \|u\|_{H^2(\Omega)}^{\frac{1}{2}}, \quad \forall u \in H^2(\Omega).$$

Furthermore, there holds, both in two and three space dimensions,

$$(4.7) \quad \|u\|_{L^\infty(\Omega)} \leq c \|u\|_{H^1(\Omega)}^{\frac{1}{2}} \|u\|_{H^2(\Omega)}^{\frac{1}{2}}, \quad \forall u \in H^2(\Omega).$$

**Remark 4.4.** Note that, interpolating as above, i.e., with exponents  $\frac{1}{2}$ , one obtains the spaces  $H^1(\Omega)$  and  $H^{\frac{3}{2}}(\Omega)$ , respectively, and that, in two space dimensions (resp., three space dimensions),  $L^\infty(\Omega)$  is not continuously embedded into  $H^1(\Omega)$  (resp.,  $H^{\frac{3}{2}}(\Omega)$ ).

4.3. **Moser–Trudinger's inequality** [340]. Let  $\Omega$  be a bounded smooth domain of  $\mathbb{R}^2$ . Then, there exists a positive constant  $C$  such that

$$(4.8) \quad \int_{\Omega} e^{|u(x)|} dx \leq C e^{C \|u\|_{H^1(\Omega)}^2}, \quad \forall u \in H^1(\Omega).$$

**Remark 4.5.** This inequality is also related to the limit case in the continuous embedding of  $H^1(\Omega)$  into  $L^p(\Omega)$  in two space dimensions and to an Orlicz space. Note that such an inequality is not available in three space dimensions.

## 5. ASYMPTOTIC BEHAVIOR OF DISSIPATIVE SYSTEMS: GLOBAL ATTRACTORS

Our aim in this section is to review some basic results in the theory of attractors for dissipative (parabolic) partial differential equations. More details and developments on this subject can be found, e.g., in [19], [335] and [388].

Let  $E$  be a Banach space endowed with the norm  $\|\cdot\|_E$ . Let then  $\{S(t), t \geq 0\}$  be a family of (nonlinear) operators acting on  $E$ ,

$$S(t) : E \rightarrow E, \quad t \geq 0.$$

We assume that this family of operators satisfies the following properties:

$$S(0) = I \text{ (identity operator),}$$

$$S(t+s) = S(t) \circ S(s), \quad \forall s, t \geq 0,$$

and we say that it forms a semigroup acting on  $E$ . We will also need some continuity property and we assume that the mapping

$$S(t) : E \rightarrow E, \quad x \mapsto S(t)x,$$

is continuous,  $\forall t \geq 0$ .

We first start with some terminology.

- For  $u_0 \in E$ , the trajectory starting at  $u_0$  (or positive trajectory) is the set  $\cup_{t \geq 0} S(t)u_0$ .
- For  $u_0 \in E$ , a trajectory ending at  $u_0$  (or negative trajectory), if it exists, is a set of the form

$$\{u(t), u(t) \in S(-t)^{-1}u_0, t \leq 0\}$$

(note that such a set, if it exists, is not necessarily unique).

- For  $u_0 \in E$ , a complete trajectory containing  $u_0$ , if it exists, is the union of the trajectory starting at  $u_0$  and of a trajectory ending at  $u_0$ . In particular, fixed points are complete trajectories ( $x \in E$  is a fixed point if  $S(t)x = x, \forall t \geq 0$ ).
- A set  $X \subset E$  is positively invariant by  $S(t)$  if  $S(t)X \subset X, \forall t \geq 0$  (thus, any trajectory starting from  $X$  remains in  $X$ ).
- A set  $X \subset E$  is negatively invariant by  $S(t)$  if  $X \subset S(t)X, \forall t \geq 0$  (thus, for every  $x \in X$ , there exists a trajectory ending at  $x$  which is contained in  $X$ ).
- A set  $X \subset E$  is invariant by  $S(t)$  if  $S(t)X = X, \forall t \geq 0$ .

For instance, complete trajectories or unions of complete trajectories are invariant by  $S(t)$ . A more complicated (and essential in view of the construction of the global attractor) invariant set is an  $\omega$ -limit set.

**Definition 5.1.** Let  $x \in E$ . The  $\omega$ -limit set of  $x$  is the set

$$\omega(x) = \cap_{s \geq 0} \overline{\cup_{t \geq s} S(t)x}.$$

Similarly, if  $B \subset E$  is a nonempty set, the  $\omega$ -limit set of  $B$  is the set

$$\omega(B) = \cap_{s \geq 0} \overline{\cup_{t \geq s} S(t)B}.$$

An important (and easy to see) characterization of  $\omega$ -limit sets is the following:

$$\begin{aligned} x \in \omega(B) \text{ iff } \exists (x_k)_{k \in \mathbb{N}}, x_k \in B, \forall k \in \mathbb{N}, \exists (t_k)_{k \in \mathbb{N}}, t_k \rightarrow +\infty \text{ as } k \rightarrow +\infty, \\ \text{such that } S(t_k)x_k \rightarrow x \text{ as } k \rightarrow +\infty. \end{aligned}$$

We have the following result (which is again essential in view of the construction of the global attractor).

**Proposition 5.2.** Let  $X \subset E$  be such that  $X \neq \emptyset$  and  $\cup_{t \geq t_0} S(t)X$  is relatively compact in  $E$  for some  $t_0 > 0$ . Then,  $\omega(X)$  is nonempty, compact and invariant.

*Proof.* First, note that, since  $X \neq \emptyset, \cup_{t \geq s} S(t)X \neq \emptyset, \forall s \geq 0$ . Therefore, the sets  $\overline{\cup_{t \geq s} S(t)X}, s \geq t_0$ , are nonempty and compact sets which decrease as  $s$  increases: their intersection, namely,  $\omega(X)$ , is a nonempty and compact set.

Let us now prove that  $\omega(X)$  is invariant by  $S(t)$ .

Let  $y \in S(t)\omega(X)$ , for a given  $t \geq 0$ . Then, there exists  $x \in \omega(X)$  such that  $y = S(t)x$ . Furthermore, owing to the characterization of  $\omega$ -limit sets, there exist  $x_k \in X$  and  $t_k \rightarrow +\infty$  such that  $S(t_k)x_k \rightarrow x$  as  $k \rightarrow +\infty$ . Then, by continuity

of  $S(t)$ ,  $S(t) \circ S(t_k)x_k \rightarrow S(t)x = y$  as  $k \rightarrow +\infty$ . Since, owing to the semigroup properties,  $S(t) \circ S(t_k)x_k = S(t + t_k)x_k$ , where  $t + t_k \rightarrow +\infty$  as  $k \rightarrow +\infty$ , it follows that  $y \in \omega(X)$  and  $S(t)\omega(X) \subset \omega(X)$ ,  $\forall t \geq 0$ .

Let now  $x \in \omega(X)$ . Then, there exists  $t_k \rightarrow +\infty$  and  $x_k \in X$  such that  $S(t_k)x_k \rightarrow x$  as  $k \rightarrow +\infty$ . Since  $\cup_{t \geq t_0} S(t)X$  is relatively compact in  $E$ , it is easy to see that the sequence  $S(t_k - t)x_k$  is relatively compact in  $E$  for  $t_k - t \geq 0$  (i.e., for  $k$  large enough). Therefore, there exists a subsequence  $k_i$  and  $y \in E$  such that

$$S(t_{k_i} - t)x_{k_i} \rightarrow y \text{ as } k_i \rightarrow +\infty.$$

This yields that  $y \in \omega(X)$  (indeed,  $t_{k_i} - t \rightarrow +\infty$  as  $k_i \rightarrow +\infty$ ). Finally, since

$$S(t) \circ S(t_{k_i} - t)x_{k_i} = S(t_{k_i})x_{k_i}$$

and, by continuity,

$$S(t) \circ S(t_{k_i} - t)x_{k_i} \rightarrow S(t)y \text{ as } k_i \rightarrow +\infty,$$

we have (of course,  $S(t_{k_i})x_{k_i} \rightarrow x$  as  $k_i \rightarrow +\infty$ )

$$x = S(t)y,$$

i.e.,  $x \in S(t)\omega(X)$ . Therefore,  $\omega(X) \subset S(t)\omega(X)$ ,  $\forall t \geq 0$ , which finishes the proof of the proposition.  $\square$

• A set  $X \subset E$  attracts the bounded sets of  $E$  (we also say that it is an attracting set) if,  $\forall B \subset E$  bounded,

$$\text{dist}_E(S(t)B, X) \rightarrow 0 \text{ as } t \rightarrow +\infty,$$

where  $\text{dist}_E$  denotes the Hausdorff semidistance between sets, defined by

$$\text{dist}_E(A, B) = \sup_{a \in A} \inf_{b \in B} \|a - b\|_E.$$

Note in particular that  $\text{dist}_E(A, B) \neq \text{dist}_E(B, A)$ . Furthermore,  $\text{dist}_E(A, B) = 0$  only implies  $A \subset \overline{B}$ . This attraction property is equivalent to the following (natural) property:

$$\forall \epsilon > 0, \forall B \subset E \text{ bounded, } \exists t_0 = t_0(B) \text{ such that } t \geq t_0 \implies S(t)B \subset \mathcal{U}_\epsilon,$$

where  $\mathcal{U}_\epsilon$  is the  $\epsilon$ -neighborhood of  $X$ , i.e., the union of all open balls with radius  $\epsilon$  and with centers in  $X$ .

• A bounded set  $\mathcal{B}_0 \subset E$  is a bounded absorbing set for  $S(t)$  if,  $\forall B \subset E$  bounded,  $\exists t_0 = t_0(B)$  such that  $t \geq t_0$  implies  $S(t)B \subset \mathcal{B}_0$ .

Note that the existence of a bounded absorbing set is sometimes used as a mathematical definition of dissipativity (from a physical point of view, roughly speaking, a dissipative system exhibits some kind of energy dissipation phenomenon).

We are now ready to give the definition of a global attractor.

**Definition 5.3.** A set  $\mathcal{A} \subset E$  is a global attractor for the semigroup  $S(t)$  if the following properties hold:

(i)  $\mathcal{A}$  is compact in  $E$ .

This essentially says that a global attractor is much thinner than the phase space  $E$ ; indeed, in applications,  $E$  usually is an infinite-dimensional function space (typically,  $L^2(\Omega)$ ,  $L^\infty(\Omega)$  or  $H_0^1(\Omega)$ ) and, in infinite dimensions, a compact set cannot contain a ball and is nowhere dense.

(ii)  $\mathcal{A}$  is invariant by  $S(t)$ ,  $S(t)\mathcal{A} = \mathcal{A}$ ,  $\forall t \geq 0$ .

(iii)  $\mathcal{A}$  is an attracting set.

We first note that a global attractor, if it exists, is unique. Indeed, if  $\mathcal{A}'$  is a second global attractor, then, since  $\mathcal{A}'$  is bounded in  $E$ ,  $\text{dist}_E(S(t)\mathcal{A}', \mathcal{A}) \rightarrow 0$  as  $t \rightarrow +\infty$  and, since, owing to the invariance,  $S(t)\mathcal{A}' = \mathcal{A}'$ , there remains  $\text{dist}_E(\mathcal{A}', \mathcal{A}) = 0$ , so that  $\mathcal{A}' \subset \overline{\mathcal{A}} = \mathcal{A}$ . Similarly,  $\mathcal{A} \subset \mathcal{A}'$  and  $\mathcal{A} = \mathcal{A}'$ .

Furthermore, it is easy to see that  $\mathcal{A}$  is the smallest (for the inclusion) closed set which enjoys the attraction property and thus appears as a suitable set in view of the study of the asymptotic behavior of the system. Indeed, let  $X \subset E$  be a closed attracting set. Then,  $\text{dist}_E(S(t)\mathcal{A}, X) \rightarrow 0$  as  $t \rightarrow +\infty$ , i.e.,  $\text{dist}_E(\mathcal{A}, X) = 0$ , so that  $\mathcal{A} \subset \overline{X} = X$ .

Finally,  $\mathcal{A}$  is the largest (again, for the inclusion) bounded invariant set. Indeed, let  $X$  be a bounded invariant set. Then,  $\text{dist}_E(S(t)X, \mathcal{A}) \rightarrow 0$  as  $t \rightarrow +\infty$ , which implies, since  $X$  is invariant and  $\mathcal{A}$  is closed, that  $X \subset \mathcal{A}$ .

**Remark 5.4.** a) The global attractor is also called the universal attractor (in the sense that it attracts all the bounded subsets of the phase space) or the maximal attractor.

b) The existence of the global attractor implies the existence of a bounded absorbing set. Indeed, any  $\epsilon$ -neighborhood of the global attractor is a bounded absorbing set.

We have, concerning the existence of the global attractor, the following result.

**Theorem 5.5.** *We assume that  $S(t)$  possesses a bounded absorbing set  $\mathcal{B}_0$  and that,  $\forall B \subset E$  bounded,  $\exists t_0 = t_0(B) > 0$  such that  $\cup_{t \geq t_0} S(t)B$  is relatively compact in  $E$ . Then,  $S(t)$  possesses the global attractor  $\mathcal{A}$  and  $\mathcal{A} = \omega(\mathcal{B}_0)$ .*

*Proof.* It actually suffices, in view of Proposition 5.2, to prove that  $\omega(\mathcal{B}_0)$  is an attracting set (indeed, we already know that  $\omega(\mathcal{B}_0)$  is a nonempty, compact and invariant set).

We will argue by contradiction.

Let us assume that  $\omega(\mathcal{B}_0)$  is not an attracting set. Then, there exists  $B_1 \subset E$  bounded such that  $\text{dist}_E(S(t)B_1, \omega(\mathcal{B}_0))$  does not tend to 0 as  $t$  tends to  $+\infty$ .

Therefore, there exist  $\delta > 0$  and  $t_k \rightarrow +\infty$  such that

$$\forall k \in \mathbb{N}, \text{dist}_E(S(t_k)B_1, \omega(\mathcal{B}_0)) \geq \delta.$$

Thus, by definition of the Hausdorff semidistance, necessarily,  $\forall k \in \mathbb{N}, \exists b_k \in B_1$  such that

$$(5.1) \quad \text{dist}_E(S(t_k)b_k, \omega(\mathcal{B}_0)) \geq \frac{\delta}{2}.$$

We recall that  $\mathcal{B}_0$  is an absorbing set. Therefore, if  $t_k \geq t_1 = t_1(B_1)$ ,  $t_1 \geq 0$ ,  $S(t_k)B_1 \subset \mathcal{B}_0$  (indeed,  $t_k \rightarrow +\infty$  as  $k \rightarrow +\infty$ ) and  $S(t_k)b_k \in \mathcal{B}_0$  if  $k$  is large enough.

Using now the fact that  $\cup_{t \geq t_0} S(t)\mathcal{B}_0$  is relatively compact for some  $t_0 > 0$ , we deduce that the sequence  $\{S(t_k)b_k, k \in \mathbb{N}\}$  is relatively compact. Therefore, there exists a subsequence  $(t_{k_i}, b_{k_i})$  and  $b \in E$  such that

$$S(t_{k_i})b_{k_i} \rightarrow b \text{ as } k_i \rightarrow +\infty.$$

Writing, for  $k_i$  large enough so that  $t_{k_i} - t_1 \geq 0$ ,

$$S(t_{k_i})b_{k_i} = S(t_{k_i} - t_1) \circ S(t_1)b_{k_i}$$

and noting that  $S(t_1)b_{k_i} \in \mathcal{B}_0$ , we deduce that  $b \in \omega(\mathcal{B}_0)$  (indeed,  $t_{k_i} - t_1 \rightarrow +\infty$  as  $k_i \rightarrow +\infty$ ). This leads to a contradiction, since, by continuity, (5.1) yields, at the limit,

$$\text{dist}_E(b, \omega(\mathcal{B}_0)) \geq \frac{\delta}{2}.$$

This finishes the proof of the theorem.  $\square$

In particular, if  $S(t)$  possesses a bounded absorbing set  $\mathcal{B}_1$  such that  $\mathcal{B}_1$  is relatively compact in  $E$ , then,  $\forall B \subset E$  bounded,  $\exists t_0 = t_0(B) > 0$  such that  $S(t)B \subset \mathcal{B}_1$ ,  $\forall t \geq t_0$ , which implies that  $\cup_{t \geq t_0} S(t)B \subset \mathcal{B}_1$  is relatively compact. Therefore, owing to Theorem 5.5,  $S(t)$  possesses the global attractor  $\mathcal{A}$ . In particular, this is the case for many parabolic systems. However, this cannot work for, e.g., weakly damped wave equations (indeed, in that case, there is no regularizing effect). In such a case, we have, for instance, the following result.

**Theorem 5.6.** *We assume that  $S(t)$  possesses a compact attracting set  $K$ . Then,  $S(t)$  possesses the global attractor  $\mathcal{A}$  and  $\mathcal{A} = \omega(K)$ .*

**Remark 5.7.** a) We can also prove, in Theorems 5.5 and 5.6, that the global attractor  $\mathcal{A}$  is connected.

b) It follows from the invariance property that the global attractor  $\mathcal{A}$  has the following structure:

$$\mathcal{A} = \{x \in E, \exists \text{ a bounded complete trajectory containing } x\}.$$

Note that any bounded complete trajectory lies in  $\mathcal{A}$  (indeed, it is a bounded invariant set).

It has been early conjectured that the global attractor should be, in some proper sense, finite-dimensional, meaning that the dynamics, restricted to this set, should be described by a finite number of parameters, even though the initial phase space is infinite-dimensional. However, the global attractor is expected to have a very complicated geometric structure (this should be compared with strange attractors in finite dimensions (i.e., for systems of ODE's in  $\mathbb{R}^n$ ,  $n \geq 3$ ) which are, typically, homeomorphic to the product of some  $\mathbb{R}^m$  and some Cantor set). Therefore, we need to use other dimensions than the usual notion of dimension to measure the dimension of the global attractor; in particular, one usually considers covering dimensions such as the Hausdorff dimension or the entropy (or box-counting) dimension. Actually, we will only consider here the entropy dimension which we will call the fractal dimension. This dimension is defined as follows.

**Definition 5.8.** Let  $X \subset E$  be a (relatively) compact set. For  $\epsilon > 0$ , let  $N_\epsilon(X)$  (if it is necessary to precise the topology, we will also use the notation  $N_\epsilon(X, E)$ ) be the minimal number of balls of radius  $\epsilon$  which are necessary to cover  $X$ . Then, the fractal dimension of  $X$  is the quantity (which belongs to  $[0, +\infty]$ )

$$\dim_{\text{F}} X = \limsup_{\epsilon \rightarrow 0^+} \frac{\log_2 N_\epsilon(X)}{\log_2 \frac{1}{\epsilon}} (= \limsup_{\epsilon \rightarrow 0^+} \frac{\ln N_\epsilon(X)}{\ln \frac{1}{\epsilon}}).$$

Furthermore, the quantity  $\mathcal{H}_\epsilon(X) (= \mathcal{H}_\epsilon(X, E)) = \log_2 N_\epsilon(X)$  is called the Kolmogorov  $\epsilon$ -entropy of  $X$ .

The fractal dimension satisfies the following properties:

- $\dim_{\text{F}}(X_1 \times X_2) \leq \dim_{\text{F}}(X_1) + \dim_{\text{F}}(X_2)$ .
- If  $f : X_1 \rightarrow X_2$  is Lipschitz continuous, then  $\dim_{\text{F}} X_2 \leq \dim_{\text{F}} X_1$ .
- If  $X$  is a smooth  $m$ -dimensional manifold, then  $\dim_{\text{F}} X = m$ .

It is also important to note that, for sets which are not manifolds, the fractal dimension can be noninteger; for instance, we have, in the case of the ternary Cantor set,

$$0 < \dim_{\text{F}} X = \frac{\ln 2}{\ln 3} < 1.$$

Furthermore, it follows from the definition that, if the minimal number of balls of radius  $\epsilon$  which are necessary to cover  $X$  satisfies

$$N_\epsilon(X) \leq c \left(\frac{1}{\epsilon}\right)^d \text{ (i.e., } \mathcal{H}_\epsilon(X) \leq d \ln \frac{1}{\epsilon} + c', \text{ } c' = \ln c),$$



where  $c$  and  $d$  are independent of  $\epsilon$ , then

$$\dim_{\mathbb{F}} X \leq d.$$

Now, an interest for considering the fractal dimension over other dimensions is given by the (modified) Hölder-Mañé theorem (see [244]). First, we need the following definition (see [245]).

**Definition 5.9.** A Borel set  $X$  of a Banach space  $E$  is prevalent if there exists a compactly supported probability measure  $\mu$  such that  $\mu(X + x) = 1$ ,  $\forall x \in E$ . A non-Borel set which contains a prevalent set is also prevalent.

We do not wish to go into details here and will just mention that prevalence extends the notion of “Lebesgue almost every” from Euclidean to infinite-dimensional spaces.

**Theorem 5.10.** (*Modified Hölder-Mañé theorem*) Assume that  $E$  is a Hilbert space and let  $X \subset E$  be a compact set such that  $\dim_{\mathbb{F}} X = d$  and  $m > 2d$  be an integer. Then, almost every (in the sense of prevalence) bounded linear projector  $P : E \rightarrow \mathbb{R}^m$  is one-to-one on  $X$  and has a Hölder continuous inverse.

It follows from this theorem that, if the global attractor  $\mathcal{A}$  has finite fractal dimension, then, fixing a projector  $P$  satisfying the assumptions of the theorem, the reduced dynamical system  $(\overline{S}(t), \overline{\mathcal{A}})$ ,  $\overline{S}(t) = P \circ S(t) \circ P^{-1}$ ,  $\overline{\mathcal{A}} = P(\mathcal{A})$ , is a finite-dimensional dynamical system (in  $\mathbb{R}^m$ ) which is Hölder continuous with respect to the initial data (note that such a result does not hold for other dimensions and, in particular, for the Hausdorff dimension which is also often considered in the theory of global attractors; see however [365] for partial results for the Hausdorff dimension). This shows that the fractal dimension plays a fundamental role in the finite-dimensional reduction of infinite-dimensional dynamical systems, although Theorem 5.10 is not completely satisfactory, since the reduced dynamical system is not Lipschitz continuous and cannot thus be realized as a well-posed system of ODE’s (indeed, one would need the Mañé projectors to be Lipschitz continuous and sufficient conditions on  $\mathcal{A}$  which would ensure that this holds are not known).

We now give a general method to prove the finite fractal dimensionality of a compact set (see [420]).

**Theorem 5.11.** Let  $X$  be a compact subset of  $E$ . We assume that there exist a Banach space  $E_1$ , with norm  $\|\cdot\|_{E_1}$ , such that  $E_1$  is compactly embedded into  $E$  and a mapping  $L : X \rightarrow X$  such that  $L(X) = X$  and

$$(5.2) \quad \|Lx_1 - Lx_2\|_{E_1} \leq c\|x_1 - x_2\|_E, \quad \forall x_1, x_2 \in X, \quad c > 0.$$

Then, the fractal dimension of  $X$  is finite and satisfies

$$(5.3) \quad \dim_{\mathbb{F}} X \leq \mathcal{H}_{\frac{1}{4c}}(B_{E_1}(0, 1), E),$$

where  $B_{E_1}(0, 1)$  is the unit ball in  $E_1$  (note that it is relatively compact in  $E$ , so that its  $\frac{1}{4c}$ -entropy is finite).

*Proof.* Let  $\{B_E(k_i, \epsilon), i = 1, \dots, N_\epsilon\}$  be an  $\epsilon$ -covering of  $X$  (note that  $X$  is compact) with balls in  $E$  with centers  $k_i \in X, i = 1, \dots, N_\epsilon$ . Then, owing to (5.2), the system

$$\{B_{E_1}(L(k_i), c\epsilon), i = 1, \dots, N_\epsilon\}$$

of balls of radius  $c\epsilon$  in  $E_1$  covers the set  $L(X)$ , i.e., it also covers  $X$ , since  $L(X) = X$ .

In a second step, we cover each of these balls with radius  $c\epsilon$  in  $E_1$  by a finite number of balls with radius  $\frac{\epsilon}{4}$  in  $E$ . By definition, the minimal number of such balls is given by

$$N_{\frac{\epsilon}{4}}(B_{E_1}(L(k_i), c\epsilon), E) = N_{\frac{\epsilon}{4}}(B_{E_1}(0, c\epsilon), E) = N_{\frac{1}{4c}}(B_{E_1}(0, 1), E) = \mathcal{N}.$$

Here, we have used the fact that  $E$  is a Banach, and thus a vector, space. If such a ball of radius  $\frac{\epsilon}{4}$  does not intersect  $X$ , there is nothing to say and we forget it. Let us now consider the balls of radius  $\frac{\epsilon}{4}$  which intersect  $X$  (and which still form a covering of  $X$ ). Note that the centers of these balls of radius  $\frac{\epsilon}{4}$  do not necessarily belong to  $X$ . Now, by considering balls of radius  $\frac{\epsilon}{2}$ , we can construct balls which still cover  $X$ , but now with centers in  $X$  (we need this to apply (5.2)), in  $E$  and with the same number of balls.

We thus deduce that

$$N_{\frac{\epsilon}{2}}(X, E) \leq \mathcal{N}N_\epsilon(X, E),$$

so that

$$(5.4) \quad \mathcal{H}_{\frac{\epsilon}{2}}(X, E) \leq \mathcal{H}_\epsilon(X, E) + \log_2 \mathcal{N}.$$

We will now prove (5.3) by proceeding recursively (i.e., by using (5.4) recursively).

First, note that, since  $X$  is compact in  $E$ , then there exist  $x_0 \in X$  and  $\epsilon_0 > 0$  such that  $X \subset B_E(x_0, \epsilon_0)$ , so that

$$(5.5) \quad \mathcal{H}_{\epsilon_0}(X, E) = 0.$$

Iterating (5.4)  $k$  times (starting from the  $\epsilon_0$ -covering by one ball) and using (5.5), we have

$$(5.6) \quad \mathcal{H}_{\frac{\epsilon_0}{2^k}}(X, E) \leq k \log_2 \mathcal{N}.$$

Let now  $\epsilon > 0$  be arbitrary and fix  $k = k(\epsilon)$  such that

$$(5.7) \quad \frac{\epsilon_0}{2^k} \leq \epsilon \leq \frac{\epsilon_0}{2^{k-1}}.$$

Then, obviously, owing to the first of (5.7),

$$N_\epsilon(X, E) \leq N_{\frac{\epsilon_0}{2^k}}(X, E),$$

so that, owing to (5.6) and the second of (5.7),

$$\mathcal{H}_\epsilon(X, E) \leq k \log_2 \mathcal{N} \leq (\log_2 \frac{\epsilon_0}{\epsilon} + 1) \log_2 \mathcal{N},$$

which yields (5.3). This finishes the proof of the theorem.  $\square$

**Remark 5.12.** a) In applications to parabolic PDE's, one usually proves (5.2) for, say,  $L = S(1)$ . Then, owing to the invariance property (i.e.,  $S(1)\mathcal{A} = \mathcal{A}$ ), one deduces that the global attractor  $\mathcal{A}$  has finite fractal dimension. For weakly damped hyperbolic systems, however, (5.2) cannot hold (we again recall that there is no regularizing effect), but one can prove weaker versions of (5.2).

b) It is also important to obtain sharp estimates on the fractal dimension of the global attractor in terms of the physical parameters of the problem. Actually, the above approach does not give the best estimates in general. Sharp estimates are usually obtained by the so-called volume contraction method, based on the Lyapunov exponents. This second approach, however, requires additional assumptions, namely, the differentiability of the semigroup, which may be difficult to prove (and may even not hold) in certain situations.

c) The existence of the finite-dimensional global attractor  $\mathcal{A}$  has been proved for a large number of important (dissipative) equations in mathematical physics, such as the two-dimensional incompressible Navier-Stokes equations, reaction-diffusion systems, weakly damped wave equations, pattern-formation equations, etc.

## REFERENCES

- [1] M. Abbaszadeh, A. Khodadadian, M. Parvizi, M. Dehghan and C. Heitzinger, *A direct meshless local collocation method for solving stochastic Cahn-Hilliard-Cook and stochastic Swift-Hohenberg equations*, *Engrg. Anal. Bound. Elem.* **98** (2019), 253-264.
- [2] H. Abels, *On a diffuse interface model for two-phase flows of viscous, incompressible fluids with matched densities*, *Arch. Ration. Mech. Anal.* **194** (2009), 463-506.
- [3] H. Abels, S. Bosia and M. Grasselli, *Cahn-Hilliard equation with nonlocal singular free energies*, *Ann. Mat. Pura Appl.* **194** (2015), 1071-1106.
- [4] H. Abels and E. Feireisl, *On a diffuse interface model for a two-phase flow of compressible viscous fluids*, *Indiana Univ. Math. J.* **57** (2008), 659-698.
- [5] H. Abels and M. Wilke, *Convergence to equilibrium for the Cahn-Hilliard equation with a logarithmic free energy*, *Nonlinear Anal.* **67** (2007), 3176-3193.
- [6] S. Agmon, *Lectures on elliptic boundary value problems*, *Mathematical Studies*, Van Nostrand, New York, 1965.
- [7] S. Agmon, A. Douglis, L. Nirenberg, *Estimates near the boundary for solutions of elliptic partial differential equations, I*, *Commun. Pure Appl. Math.* **12** (1959), 623-727.
- [8] S. Agmon, A. Douglis, L. Nirenberg, *Estimates near the boundary for solutions of elliptic partial differential equations, II*, *Commun. Pure Appl. Math.* **17** (1964), 35-92.
- [9] M. Ainsworth and Z. Mao, *Well-posedness of the Cahn-Hilliard equation with fractional free energy and its Fourier Galerkin approximation*, *Chaos Solitons Fractals* **102** (2017), 264-273.

- [10] M. Ainsworth and Z. Mao, *Analysis and approximation of a fractional Cahn-Hilliard equation*, SIAM J. Numer. Anal. **55** (2017), 1689-1718.
- [11] S.M. Allen and J.W. Cahn, *A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening*, Acta Metall. **27** (1979), 1085-1095.
- [12] H.W. Alt and I. Pawlow, *A mathematical model of dynamics of non-isothermal phase separation*, Phys. D **59** (1992), 389-416.
- [13] H.W. Alt and I. Pawlow, *Existence of solutions for non-isothermal phase separation*, Adv. Math. Sci. Appl. **1** (1992), 319-409.
- [14] P.F. Antonietti, L. Beirão da Veiga, S. Scacchi and M. Verani,  *$C^1$  virtual element method for the Cahn-Hilliard equation with polygonal meshes*, SIAM J. Numer. Anal. **54** (2016), 34-56.
- [15] A.R. Appadu, J.K. Djoko, H.H. Gidey and J.M.S. Lubuma, *Analysis of multilevel finite volume approximation of 2D convective Cahn-Hilliard equation*, Jpn. J. Ind. Appl. Math. **34** (2017), 253-304.
- [16] P. Areias, E. Samaniego and T. Rabczuk, *A staggered approach for the coupling of Cahn-Hilliard type diffusion and finite strain elasticity*, Comput. Mech. **57** (2016), 339-351.
- [17] A. Aristotelous, O. Karakashian and S.M. Wise, *A mixed discontinuous Galerkin, convex splitting scheme for a modified Cahn-Hilliard equation and an efficient nonlinear multigrid solver*, Discrete Contin. Dyn. Systems Ser. B **18** (2013), 2211-2238.
- [18] A.C. Aristotelous, O.A. Karakashian and S.M. Wise, *Adaptive, second-order in time, primitive-variable discontinuous Galerkin schemes for a Cahn-Hilliard equation with a mass source*, IMA J. Numer. Anal. **35** (2015), 1167-1198.
- [19] A.V. Babin and M.I. Vishik, *Attractors of evolution equations*, North-Holland, Amsterdam, 1992.
- [20] F. Bai, C.M. Elliott, A. Gardiner, A. Spence and A.M. Stuart, *The viscous Cahn-Hilliard equation. Part I: computations*, Nonlinearity **8** (1995), 131-160.
- [21] L. Bañas and R. Nürnberg, *A multigrid method for the Cahn-Hilliard equation with obstacle potential*, Appl. Math. Comput. **213** (2009), 290-303.
- [22] J.W. Barrett and J.F. Blowey, *An error bound for the finite element approximation of the Cahn-Hilliard equation with logarithmic free energy*, Numer. Math. **72** (1995), 257-287.
- [23] J.W. Barrett and J.F. Blowey, *Finite element approximation of the Cahn-Hilliard equation with concentration dependent mobility*, Math. Comput. **68** (1999), 487-517.
- [24] J.W. Barrett, J.F. Blowey and H. Garcke, *Finite element approximation of the Cahn-Hilliard equation with degenerate mobility*, SIAM J. Numer. Anal. **37** (1999), 286-318.
- [25] S. Bartels and R. Müller, *A posteriori error controlled local resolution of evolving interfaces for generalized Cahn-Hilliard equations*, Interfaces Free Bound. **12** (2010), 45-73.
- [26] S. Bartels and R. Müller, *Error control for the approximation of Allen-Cahn and Cahn-Hilliard equations with a logarithmic potential*, Numer. Math. **119** (2011), 409-435.
- [27] P.W. Bates and J. Han, *The Neumann boundary problem for a nonlocal Cahn-Hilliard equation*, J. Diff. Eqns. **212** (2005), 235-277.
- [28] A. Bertozzi, S. Esedoglu and A. Gillette, *Inpainting of binary images using the Cahn-Hilliard equation*, IEEE Trans. Imag. Proc. **16** (2007), 285-291.
- [29] A. Bertozzi, S. Esedoglu and A. Gillette, *Analysis of a two-scale Cahn-Hilliard model for binary image inpainting*, Multiscale Model. Simul. **6** (2007), 913-936.
- [30] K. Binder and H.L. Frisch, *Dynamics of surface enrichment: A theory based on the Kawasaki spin-exchange model in the presence of a wall*, Z. Phys. B **84** (1991), 403-418.

- [31] T. Blesgen and I.V. Chenchiah, *Cahn-Hilliard equations incorporating elasticity: analysis and comparison to experiments*, Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Engrg. Sci. **371** (2013), 1-16.
- [32] T. Blesgen and A. Schlömerkemper, *On the Allen-Cahn/Cahn-Hilliard system with a geometrically linear elastic energy*, Proc. Roy. Soc. Edinburgh Sect. A **144** (2014), 241-266.
- [33] D. Blömker, B. Gawron and T. Wanner, *Nucleation in the one-dimensional stochastic Cahn-Hilliard model*, Discrete Contin. Dyn. Systems **27** (2010), 25-52.
- [34] D. Blömker, S. Maier-Paape and T. Wanner, *Second phase spinodal decomposition for the Cahn-Hilliard-Cook equation*, Trans. Amer. Math. Soc. **360** (2008), 449-489.
- [35] J.F. Blowey and C.M. Elliott, *The Cahn-Hilliard gradient theory for phase separation with non-smooth free energy. Part I. Mathematical analysis*, Eur. J. Appl. Math. **2** (1991), 233-280.
- [36] J.F. Blowey and C.M. Elliott, *The Cahn-Hilliard gradient theory for phase separation with non-smooth free energy. Part II. Numerical analysis*, Eur. J. Appl. Math. **3** (1992), 147-179.
- [37] L. Bo, K. Shi and Y. Wang, *Support theorem for a stochastic Cahn-Hilliard equation*, Electron. J. Prob. **15** (2010), 484-525.
- [38] L. Bo and Y. Wang, *Stochastic Cahn-Hilliard partial differential equations with Lévy spacetime white noise*, Stochastics Dyn. **6** (2006), 229-244.
- [39] E. Bonetti, P. Colli, W. Dreyer, G. Gilardi, G. Schimperna and J. Sprekels, *On a model for phase separation in binary alloys driven by mechanical effects*, Phys. D **165** (2002), 48-65.
- [40] E. Bonetti, W. Dreyer and G. Schimperna, *Global solution to a viscous Cahn-Hilliard equation for tin-lead alloys with mechanical stresses*, Adv. Diff. Eqns. **2** (2003), 231-256.
- [41] A. Bonfoh, M. Grasselli and A. Miranville, *Long time behavior of a singular perturbation of the viscous Cahn-Hilliard-Gurtin equation*, Math. Methods Appl. Sci. **31** (2008), 695-734.
- [42] A. Bonfoh, M. Grasselli and A. Miranville, *Inertial manifolds for a singular perturbation of the viscous Cahn-Hilliard-Gurtin equation*, Topol. Methods Nonlinear Anal. **35** (2010), 155-185.
- [43] A. Bonfoh, M. Grasselli and A. Miranville, *Singularly perturbed 1D Cahn-Hilliard equation revisited*, Nonlinear Diff. Eqns. Appl. (NoDEA) **17** (2010), 663-695.
- [44] A. Bonfoh and A. Miranville, *On Cahn-Hilliard-Gurtin equations*, Nonlinear Anal. **47** (2001), 3455-3466.
- [45] J. Bosch and M. Stoll, *Preconditioning for vector-valued Cahn-Hilliard equations*, SIAM J. Sci. Comput. **37** (2015), S216-S243.
- [46] J. Bosch and M. Stoll, *A fractional inpainting model based on the vector-valued Cahn-Hilliard equation*, SIAM J. Imag. Sci. **8** (2015), 2352-2382.
- [47] S. Bosia, M. Conti and M. Grasselli, *On the Cahn-Hilliard-Brinkman system*, Commun. Math. Sci. **13** (2015), 1541-1567.
- [48] S. Bosia, M. Grasselli and A. Miranville, *On the longtime behavior of a 2D hydrodynamic model for chemically reacting binary fluid mixtures*, Math. Methods Appl. Sci. **37** (2014), 726-743.
- [49] F. Boyer, *Nonhomogeneous Cahn-Hilliard fluids*, Ann. Inst. H. Poincaré Anal. Non Linéaire **18** (2001), 225-259.
- [50] F. Boyer, *A theoretical and numerical model for the study of incompressible mixture flows*, Computers and Fluids **31** (2002), 41-68.
- [51] F. Boyer, L. Chupin and P. Fabrie, *Numerical study of viscoelastic mixtures through a Cahn-Hilliard flow model*, Eur. J. Mech. B Fluids **23** (2004), 759-780.
- [52] F. Boyer and C. Lapuerta, *Study of a three component Cahn-Hilliard flow model*, M2AN Math. Model. Numer. Anal. **40** (2006), 653-687.

- [53] F. Boyer, C. Lapuerta, S. Minjeaud and B. Piar, *A local adaptive refinement method with multigrid preconditioning illustrated by multiphase flows simulations*, ESAIM Proc. **27** (2009), 15-53.
- [54] F. Boyer, C. Lapuerta, S. Minjeaud, B. Piar and M. Quintard, *Cahn-Hilliard/Navier-Stokes model for the simulation of three-phase flows*, Transp. Porous Media **82** (2010), 463-483.
- [55] F. Boyer and S. Minjeaud, *Numerical schemes for a three component Cahn-Hilliard model*, M2AN Math. Model. Numer. Anal. **45** (2011), 697-738.
- [56] F. Boyer and S. Minjeaud, *Hierarchy of consistent n-component Cahn-Hilliard systems*, Math. Models Methods Appl. Sci. **24** (2014), 2885-2928.
- [57] F. Boyer and F. Nabet, *A DDFV method for a Cahn-Hilliard/Stokes phase field model with dynamic boundary conditions*, ESAIM Math. Model. Numer. Anal. **51** (2017), 1691-1731.
- [58] S.C. Brenner, A.E. Diegel and L.-Y. Sung, *A robust solver for a mixed finite element method for the Cahn-Hilliard equation*, J. Sci. Comput. **77** (2018), 1234-1249.
- [59] L.A. Caffarelli and N.E. Muler, *An  $L^\infty$  bound for solutions of the Cahn-Hilliard equation*, Arch. Ration. Mech. Anal. **133** (1995), 129-144.
- [60] G. Caginalp and E. Esenturk, *Anisotropic phase field equations of arbitrary order*, Discrete Contin. Dyn. Systems Ser. S **4** (2011), 311-350.
- [61] J.W. Cahn, *On spinodal decomposition*, Acta Metall. **9** (1961), 795-801.
- [62] J.W. Cahn, C.M. Elliott and A. Novick-Cohen, *The Cahn-Hilliard equation with a concentration dependent mobility: motion by minus the Laplacian of the mean curvature*, Eur. J. Appl. Math. **7** (1996), 287-301.
- [63] J.W. Cahn and J.E. Hilliard, *Free energy of a nonuniform system I. Interfacial free energy*, J. Chem. Phys. **28** (1958), 258-267.
- [64] J.W. Cahn and J.E. Hilliard, *Free energy of a nonuniform system III. Nucleation in a two component incompressible fluid*, J. Chem. Phys. **31** (1959), 688-699.
- [65] Y. Cai, H. Choi and J. Shen, *Error estimates for time discretizations of Cahn-Hilliard and Allen-Cahn phase-field models for two-phase incompressible flows*, Numer. Math. **137** (2017), 417-449.
- [66] Y. Cai and J. Shen, *Error estimates for a fully discretized scheme to a Cahn-Hilliard phase-field model for two-phase incompressible flows*, Math. Comp. **87** (2018), 2057-2090.
- [67] C. Cao and C.G. Gal, *Global solutions for the 2D NS-CH model for a two-phase flow of viscous, incompressible fluids with mixed partial viscosity and mobility*, Nonlinearity **25** (2012), 3211-3234.
- [68] C. Cardon-Weber, *Cahn-Hilliard stochastic equation: existence of the solution and of its density*, Bernoulli **7** (2001), 777-816.
- [69] M. Carrive, A. Miranville and A. Piétrus, *The Cahn-Hilliard equation for deformable continua*, Adv. Math. Sci. Appl. **10** (2000), 539-569.
- [70] A.N. Carvalho and T. Dlotko, *Dynamics of the viscous Cahn-Hilliard equation*, J. Math. Anal. Appl. **344** (2008), 703-725.
- [71] S. Chai, Y. Cao, Y. Zou and W. Zhao, *Conforming finite element methods for the stochastic Cahn-Hilliard-Cook equation*, Appl. Numer. Math. **124** (2018), 44-56.
- [72] V. Chalupeckí, *Numerical studies of Cahn-Hilliard equations and applications in image processing*, in Proceedings of Czech-Japanese Seminar in Applied Mathematics 2004 (August 4-7, 2004), Czech Technical University in Prague.
- [73] F. Chave, D.A. Di Pietro, F. Marche and F. Pigeonneau, *A hybrid high-order method for the Cahn-Hilliard problem in mixed form*, SIAM J. Numer. Anal. **54** (2016), 1873-1898.

- [74] F. Chen and J. Shen, *Efficient energy stable schemes with spectral discretization in space for anisotropic Cahn-Hilliard systems*, Commun. Comput. Phys. **13** (2013), 1189-1208.
- [75] L. Chen and C. Xu, *A time splitting space spectral element method for the Cahn-Hilliard equation*, East Asian J. Appl. Math. **3** (2013), 333-351.
- [76] X. Chen, G. Caginalp and E. Esenturk, *Interface conditions for a phase field model with anisotropic and non-local interactions*, Arch. Rational Mech. Anal. **202** (2011), 349-372.
- [77] K. Cheng, C. Wang, S.M. Wise and Y. Yue, *A second-order, weakly energy-stable pseudo-spectral scheme for the Cahn-Hilliard equation and its solution by the homogeneous linear iteration method*, J. Sci. Comput. **69** (2016), 1083-1114.
- [78] L. Cherfils, H. Fakhir, M. Grasselli and A. Miranville, *A convergent convex splitting scheme for a nonlocal Cahn-Hilliard-Oono type equation with a transport term*, submitted.
- [79] L. Cherfils, H. Fakhir and A. Miranville, *Finite-dimensional attractors for the Bertozzi-Esedoglu-Gillette-Cahn-Hilliard equation in image inpainting*, Inv. Prob. Imag. **9** (2015), 105-125.
- [80] L. Cherfils, H. Fakhir and A. Miranville, *On the Bertozzi-Esedoglu-Gillette-Cahn-Hilliard equation with logarithmic nonlinear terms*, SIAM J. Imag. Sci. **8** (2015), 1123-1140.
- [81] L. Cherfils, H. Fakhir and A. Miranville, *A Cahn-Hilliard system with a fidelity term for color image inpainting*, J. Math. Imag. Vision **54** (2016), 117-131.
- [82] L. Cherfils, H. Fakhir and A. Miranville, *A complex version of the Cahn-Hilliard equation for grayscale image inpainting*, Multiscale Model. Simul. **15** (2017), 575-605.
- [83] L. Cherfils, S. Gatti and Miranville, *A variational approach to a Cahn-Hilliard model in a domain with nonpermeable walls*, J. Math. Sci. (N.Y.) **189** (2012), 604-636.
- [84] L. Cherfils, S. Gatti and Miranville, *Asymptotic behavior of higher-order Navier-Stokes-Cahn-Hilliard systems*, Math. Methods Appl. Sci. **41** (2018), 4776-4794.
- [85] L. Cherfils and A. Miranville, *Generalized Cahn-Hilliard equations with a logarithmic free energy*, Rev. Real Acad. Sci. **94** (2000), 19-32.
- [86] L. Cherfils, A. Miranville and S. Peng, *Higher-order models in phase separation*, J. Appl. Anal. Comput. **7** (2017), 39-56.
- [87] L. Cherfils, A. Miranville and S. Peng, *Higher-order anisotropic models in phase separation*, Adv. Nonlinear Anal., to appear.
- [88] L. Cherfils, A. Miranville, S. Peng and W. Zhang, *Higher-order generalized Cahn-Hilliard equations*, Electron. J. Qualitative Theory Diff. Eqns. **9** (2017), 1-22.
- [89] L. Cherfils, A. Miranville and S. Zelik, *The Cahn-Hilliard equation with logarithmic potentials*, Milan J. Math. **79** (2011), 561-596.
- [90] L. Cherfils, A. Miranville and S. Zelik, *On a generalized Cahn-Hilliard equation with biological applications*, Discrete Contin. Dyn. Systems Ser. B **19** (2014), 2013-2026.
- [91] L. Cherfils and M. Petcu, *On the viscous Cahn-Hilliard-Navier-Stokes equations with dynamic boundary conditions*, Commun. Pure Appl. Anal. **15** (2016), 1419-1449.
- [92] L. Cherfils, M. Petcu and M. Pierre, *A numerical analysis of the Cahn-Hilliard equation with dynamic boundary conditions*, Discrete Contin. Dyn. Systems **27** (2010), 1511-1533.
- [93] R. Chill, E. Fařangova and J. Pruss, *Convergence to steady states of solutions of the Cahn-Hilliard equation with dynamic boundary conditions*, Math. Nachr. **279** (2006), 1448-1462.
- [94] Y. Choi, D. Jeong and J. Kim, *A multigrid solution for the Cahn-Hilliard equation on nonuniform grids*, Appl. Math. Comput. **293** (2017), 320-333.
- [95] R. Choksi and X. Ren, *On a derivation of a density functional theory for microphase separation of diblock copolymers*, J. Statist. Phys. **113** (2003), 151-176.

- [96] J.W. Cholewa and T. Dlotko, *Global attractor for the Cahn-Hilliard system*, Bull. Austral. Math. Soc. **49** (1994), 277-293.
- [97] J.W. Cholewa and T. Dlotko, *Global attractors in abstract parabolic problems*, London Mathematical Society Lecture Notes Series, Vol. **278**, Cambridge University Press, Cambridge, 2000.
- [98] S.M. Choo and S.K. Chung, *Asymptotic behaviour of the viscous Cahn-Hilliard equation*, J. Appl. Math. Comput. **11** (2003), 143-154.
- [99] L. Chupin, *An existence result for a mixture of non-newtonian fluids with stress-diffusion using the Cahn-Hilliard formulation*, Discrete Contin. Dyn. Systems Ser. B **3** (2003), 45-68.
- [100] D. Cohen and J.M. Murray, *A generalized diffusion model for growth and dispersion in a population*, J. Math. Biol. **12** (1981), 237-248.
- [101] P. Colli, G. Gilardi and D. Hillhorst, *On a Cahn-Hilliard type phase field model related to tumor growth*, Discrete Contin. Dyn. Systems **35** (2015), 2423-2442.
- [102] P. Colli, G. Gilardi, P. Krejci and J. Sprekels, *A continuous dependence result for a nonstandard system of phase field equations*, Math. Methods Appl. Sci. **37** (2014), 1318-1324.
- [103] P. Colli, G. Gilardi, P. Podio-Guidugli and J. Sprekels, *Global existence for a strongly coupled Cahn-Hilliard system with viscosity*, Boll. Unione Mat. Ital. **5** (2012), 495-513.
- [104] P. Colli, G. Gilardi, P. Podio-Guidugli and J. Sprekels, *An asymptotic analysis for a nonstandard Cahn-Hilliard system with viscosity*, Discrete Contin. Dyn. Systems Ser. S **6** (2013), 353-368.
- [105] P. Colli, G. Gilardi, P. Podio-Guidugli and J. Sprekels, *Global existence and uniqueness for a singular/degenerate Cahn-Hilliard system with viscosity*, J. Diff. Eqns. **254** (2013), 4217-4244.
- [106] P. Colli, G. Gilardi, E. Rocca and J. Sprekels, *Vanishing viscosities and error estimate for a Cahn-Hilliard type phase-field system related to tumor growth*, Nonlinear Anal. Real World Appl. **26** (2015), 93-108.
- [107] P. Colli, G. Gilardi, E. Rocca and J. Sprekels, *Asymptotic analyses and error estimates for a Cahn-Hilliard type phase field system modelling tumor growth*, Discrete Contin. Dyn. Systems Ser. S **10** (2017), 37-54.
- [108] P. Colli, G. Gilardi, E. Rocca and J. Sprekels, *Optimal distributed control of a diffuse interface model of tumor growth*, Nonlinearity **30** (2017), 2518-2546.
- [109] P. Colli, G. Gilardi and J. Sprekels, *On the Cahn-Hilliard equation with dynamic boundary conditions and a dominating boundary potential*, J. Math. Anal. Appl. **419** (2014), 972-994.
- [110] P. Colli, G. Gilardi and J. Sprekels, *Global existence for a nonstandard viscous Cahn-Hilliard system with dynamic boundary condition*, SIAM J. Math. Anal. **49** (2017), 1732-1760.
- [111] M. Conti and M. Coti Zelati, *Attractors for the Cahn-Hilliard equation with memory in 2D*, Nonlinear Anal. **72** (2010), 1668-1682.
- [112] M. Conti, S. Gatti and A. Miranville, *Multi-component Cahn-Hilliard systems with dynamic boundary conditions*, Nonlinear Anal. Real World Appl. **25** (2015), 137-166.
- [113] M. Conti, S. Gatti and A. Miranville, *A singular Cahn-Hilliard-Oono phase-field system with hereditary memory*, Discrete Contin. Dynam. Systems **38** (2018), 3033-3054.
- [114] M. Conti and A. Giorgini, *On the Cahn-Hilliard-Brinkman system with singular potential and nonconstant viscosity*, submitted.
- [115] M. Conti and G. Mola, *3-D viscous Cahn-Hilliard equation with memory*, Math. Methods Appl. Sci. **32** (2009), 1370-1395.
- [116] H. Cook, *Brownian motion in spinodal decomposition*, Acta Metall. **18** (1970), 297-306.
- [117] M.I.M. Copetti and C.M. Elliott, *Numerical analysis of the Cahn-Hilliard equation with a logarithmic free energy*, Numer. Math. **63** (1992), 39-65.



- [118] G. Da Prato and A. Debussche, *Stochastic Cahn-Hilliard equation*, *Nonlinear Anal.* **26** (1996), 241-263.
- [119] M. Dai, E. Feireisl, E. Rocca, G. Schimperna and M.E. Schonbek, *Analysis of a diffuse interface model of multispecies tumor growth*, *Nonlinearity* **30** (2017), 1639-1658.
- [120] R. Dal Passo, L. Giacomelli and A. Novick-Cohen, *Existence for an Allen-Cahn/Cahn-Hilliard system with degenerate mobility*, *Interfaces Free Bound.* **1** (1999), 199-226.
- [121] P.G. de Gennes, *Dynamics of fluctuations and spinodal decomposition in polymer blends*, *J. Chem. Phys.* **72** (1980), 4756-4763.
- [122] A. Debussche and L. Dettori, *On the Cahn-Hilliard equation with a logarithmic free energy*, *Nonlinear Anal.* **24** (1995), 1491-1514.
- [123] A. Debussche and L. Goudenège, *Stochastic Cahn-Hilliard equation with double singular nonlinearities and two reflections*, *SIAM J. Math. Anal.* **43** (2011), 1473-1494.
- [124] A. Debussche and L. Zambotti, *Conservative stochastic Cahn-Hilliard equation with reflection*, *Ann. Probab.* **35** (2007), 1706-1739.
- [125] L. Dede, H. Garcke and K.F. Lam, *A Hele-Shaw-Cahn-Hilliard model for incompressible two-phase flows with different densities* *J. Math. Fluid Mech.* **20** (2018), 531-567.
- [126] F. Della Porta, A. Giorgini and M. Grasselli, *The nonlocal Cahn-Hilliard-Hele-Shaw system with singular potential*, submitted.
- [127] F. Della Porta and M. Grasselli, *On the nonlocal Cahn-Hilliard-Brinkman and Cahn-Hilliard-Hele-Shaw systems*, *Commun. Pure Appl. Anal.* **15** (2016), 299-317. Erratum, *Commun. Pure Appl. Anal.* **16** (2017), 369-372.
- [128] C.V. Di Leo, E. Rejovitzky and L. Anand, *A Cahn-Hilliard-type phase-field theory for species diffusion coupled with large elastic deformations: application to phase-separating Li-ion electrode materials*, *J. Mech. Phys. Solids* **70** (2014), 1-29.
- [129] A.E. Diegel, C. Wang, X. Wang and S.M. Wise, *Convergence analysis and error estimates for a second order accurate finite element method for the Cahn-Hilliard-Navier-Stokes system*, *Numer. Math.* **137** (2017), 495-534.
- [130] A. Diegel, C. Wang and S.M. Wise, *Stability and convergence of a second order mixed finite element method for the Cahn-Hilliard equation*, *IMA J. Numer. Anal.* **36** (2016), 1867-1897.
- [131] T. Dlotko, *Global attractor for the Cahn-Hilliard equation in  $H^2$  and  $H^3$* , *J. Diff. Eqns.* **113** (1994), 381-393.
- [132] I.C. Dolcetta and S.F. Vita, *Area-preserving curve-shortening flows: from phase separation to image processing*, *Interfaces Free Bound.* **4** (2002), 325-343.
- [133] Q. Du, L. Ju, X. Li and Z. Qiao, *Stabilized linear semi-implicit schemes for the nonlocal Cahn-Hilliard equation*, *J. Comput. Phys.* **363** (2018), 39-54.
- [134] A. Eden, C. Foias, B. Nicolaenko and R. Temam, *Exponential attractors for dissipative evolution equations*, *Research in Applied Mathematics*, Vol. **37**, John-Wiley, New York, 1994.
- [135] A. Eden and V.K. Kalantarov, *The convective Cahn-Hilliard equation*, *Appl. Math. Lett.* **20** (2007), 455-461.
- [136] A. Eden and V.K. Kalantarov, *3D convective Cahn-Hilliard equation*, *Commun. Pure Appl. Anal.* **6** (2007), 1075-1086.
- [137] A. Eden, V. Kalantarov and S.V. Zelik, *Global solvability and blow up for the convective Cahn-Hilliard equations with concave potentials*, *J. Math. Phys.* **54** (2013), 1-12.
- [138] A. Eden, V. Kalantarov and S.V. Zelik, *Infinite energy solutions for the Cahn-Hilliard equation in cylindrical domains*, *Math. Methods Appl. Sci.* **37** (2014), 1884-1908.

- [139] M. Efendiev, H. Gajewski and S. Zelik, *The finite dimensional attractor for a 4th order system of the Cahn-Hilliard type with a supercritical nonlinearity*, Adv. Diff. Eqns. **7** (2002), 1073-1100.
- [140] M. Efendiev and A. Miranville, *New models of Cahn-Hilliard-Gurtin equations*, Contin. Mech. Thermodyn. **16** (2004), 441-451.
- [141] M. Efendiev, A. Miranville and S. Zelik, *Exponential attractors for a singularly perturbed Cahn-Hilliard system*, Math. Nach. **272** (2004), 11-31.
- [142] N. Elezovic and A. Mikelic, *On the stochastic Cahn-Hilliard equation*, Nonlinear Anal. **16** (1991), 1169-1200.
- [143] C.M. Elliott, *The Cahn-Hilliard model for the kinetics of phase separation*, in Mathematical models for phase change problems, J.F. Rodrigues ed., International Series of Numerical Mathematics, Vol. **88**, Birkhäuser, Basel, 1989.
- [144] C.M. Elliott and D.A. French, *Numerical studies of the Cahn-Hilliard equation for phase separation*, IMA J. Appl. Math. **38** (1987), 97-128.
- [145] C.M. Elliott and D.A. French, *A non-conforming finite element method for the two-dimensional Cahn-Hilliard equation*, SIAM J. Numer. Anal. **26** (1989), 884-903.
- [146] C.M. Elliott, D.A. French and F.A. Milner, *A second order splitting method for the Cahn-Hilliard equation*, Numer. Math. **54** (1989), 575-590.
- [147] C.M. Elliott and H. Garcke, *On the Cahn-Hilliard equation with degenerate mobility*, SIAM J. Math. Anal. **27** (1996), 404-423.
- [148] C.M. Elliott and H. Garcke, *Diffusional phase transitions in multicomponent systems with a concentration dependent mobility matrix*, Phys. D **109** (1997), 242-256.
- [149] C.M. Elliott and S. Luckhaus, *A generalized diffusion equation for phase separation of a multi-component mixture with interfacial energy*, SFB 256 Preprint No. **195**, University of Bonn, 1991.
- [150] C.M. Elliott and T. Ranner, *Evolving surface finite element method for the Cahn-Hilliard equation*, Numer. Math. **129** (2015), 483-534.
- [151] C.M. Elliott and A.M. Stuart, *Viscous Cahn-Hilliard equation II. Analysis*, J. Diff. Eqns. **128** (1996), 387-414.
- [152] C.M. Elliott and S. Zheng, *On the Cahn-Hilliard equation*, Arch. Ration. Mech. Anal. **96** (1986), 339-357.
- [153] H. Emmerich, H. Löwen, R. Wittkowski, T. Gruhn, G.I. Tóth, G. Tegze and L. Gránásy, *Phase-field-crystal models for condensed matter dynamics on atomic length and diffusive time scales: an overview*, Adv. Phys. **61** (2012), 665-743.
- [154] J. Erlebacher, M.J. Aziz, A. Karma, N. Dimitrov and K. Sieradzki, *Evolution of nanoporosity in dealloying*, Nature **410** (2001), 450-453.
- [155] J.D. Eyre, *Systems of Cahn-Hilliard equations*, SIAM J. Appl. Math. **53** (1993), 1686-1712.
- [156] J.D. Eyre, *Unconditionally gradient stable time marching the Cahn-Hilliard equation*, in Computational and mathematical models of microstructural evolution, J.W. Bullard, R. Kalia, M. Stoneham and L.Q. Chen eds., The Materials Research Society, 1998.
- [157] H. Fakhir, *A Cahn-Hilliard equation with a proliferation term for biological and chemical applications*, Asymptot. Anal. **94** (2015), 71-104.
- [158] H. Fakhir, *Asymptotic behavior of a generalized Cahn-Hilliard equation with a mass source*, Appl. Anal. **96** (2017), 324-348.
- [159] X. Feng, Y. Li and Y. Xing, *Analysis of mixed interior penalty discontinuous Galerkin methods for the Cahn-Hilliard equation and the Hele-Shaw flow*, SIAM J. Numer. Anal. **54** (2016), 825-847.

- [160] X. Feng and A. Prohl, *Error analysis of a mixed finite element method for the Cahn-Hilliard equation*, Numer. Math. **99** (2004), 47-84.
- [161] X. Feng and A. Prohl, *Numerical analysis of the Cahn-Hilliard equation and approximation for the Hele-Shaw problem*, Interfaces Free bound. **7** (2005), 1-28.
- [162] X. Feng and S. Wise, *Analysis of a Darcy-Cahn-Hilliard diffuse interface model for the Hele-Shaw flow and its fully discrete finite element approximation*, SIAM J. Numer. Anal. **50** (2012), 1320-1343.
- [163] W.M. Feng, P. Yu, S.Y. Hu, Z.K. Liu, Q. Du and L.Q. Chen, *A Fourier spectral moving mesh method for the Cahn-Hilliard equation with elasticity*, Commun. Comput. Phys. **5** (2009), 582-599.
- [164] H.P. Fischer, P. Maass and W. Dieterich, *Novel surface modes in spinodal decomposition*, Phys. Rev. Lett. **79** (1997), 893-896.
- [165] H.P. Fischer, P. Maass and W. Dieterich, *Diverging time and length scales of spinodal decomposition modes in thin flows*, Europhys. Lett. **42** (1998), 49-54.
- [166] H.P. Fischer, J. Reinhard, W. Dieterich, J.-F. Gouyet, P. Maass, A. Majhofer and D. Reinel, *Time-dependent density functional theory and the kinetics of lattice gas systems in contact with a wall*, J. Chem. Phys. **108** (1998), 3028-3037.
- [167] S. Frigeri and M. Grasselli, *Nonlocal Cahn-Hilliard-Navier-Stokes systems with singular potentials*, Dyn. Partial Diff. Eqns. **9** (2012), 273-304.
- [168] S. Frigeri, M. Grasselli and D. Prazak, *Nonlocal Cahn-Hilliard-Navier-Stokes systems with shear dependent viscosity*, J. Math. Anal. Appl. **459** (2018), 753-777.
- [169] S. Frigeri, M. Grasselli and E. Rocca, *On a diffuse interface model of tumor growth*, Eur. J. Appl. Math. **26** (2015), 215-243.
- [170] T. Fukao, S. Yoshikawa and S. Wada, *Structure-preserving finite difference schemes for the Cahn-Hilliard equation with dynamic boundary conditions in the one-dimensional case*, Commun. Pure Appl. Anal. **16** (2017), 1915-1938.
- [171] D. Furihata, *A stable and conservative finite difference scheme for the Cahn-Hilliard equation*, Numer. Math. **87** (2001), 675-699.
- [172] D. Furihata, M. Kovacs, S. Larsson and F. Lindgren, *Strong convergence of a fully discrete finite element approximation of the stochastic Cahn-Hilliard equation*, SIAM J. Numer. Anal. **56** (2018), 708-731.
- [173] H. Gajewski and K. Zacharias, *On a nonlocal phase separation model*, J. Math. Anal. Appl. **286** (2003), 11-31.
- [174] C.G. Gal, *A Cahn-Hilliard model in bounded domains with permeable walls*, Math. Methods Appl. Sci. **29** (2006), 2009-2036.
- [175] C.G. Gal, *Exponential attractors for a Cahn-Hilliard model in bounded domains with permeable walls*, Electron. J. Diff. Eqns. **2006** (2006), 1-23.
- [176] C.G. Gal, *Non-local Cahn-Hilliard equations with fractional dynamic boundary conditions*, Eur. J. Appl. Math. **28** (2017), 736-788.
- [177] C.G. Gal, *Doubly nonlocal Cahn-Hilliard equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire **35** (2018), 357-392.
- [178] C.G. Gal, A. Giorgini and M. Grasselli, *The nonlocal Cahn-Hilliard equation with singular potential: well-posedness, regularity and strict separation property*, J. Diff. Eqns. **263** (2017), 5253-5297.
- [179] C.G. Gal and M. Grasselli, *Asymptotic behavior of a Cahn-Hilliard-Navier-Stokes system in 2D*, Ann. Inst. H. Poincaré Anal. Non Linéaire **27** (2010), 401-436.

- [180] C.G. Gal and M. Grasselli, *Longtime behavior for a model of homogeneous incompressible two-phase flows*, Discrete Contin. Dyn. Systems **28** (2010), 1-39.
- [181] C.G. Gal and M. Grasselli, *Longtime behavior of nonlocal Cahn-Hilliard equations*, Discrete Contin. Dyn. Systems **34** (2014), 145-179.
- [182] C.G. Gal, M. Grasselli and A. Miranville, *Cahn-Hilliard-Navier-Stokes systems with moving contact lines*, Calc. Var. Partial Diff. Eqns. **55** (2016), 1-47.
- [183] C.G. Gal and A. Miranville, *Uniform global attractors for non-isothermal viscous and non-viscous Cahn-Hilliard equations with dynamic boundary conditions*, Nonlinear Anal. Real World Appl. **10** (2009), 1738-1766.
- [184] C.G. Gal and A. Miranville, *Robust exponential attractors and convergence to equilibria for non-isothermal Cahn-Hilliard equations with dynamic boundary conditions*, Discrete Contin. Dyn. Systems Ser. S **2** (2009), 113-147.
- [185] C.G. Gal and H. Wu, *Asymptotic behavior of a Cahn-Hilliard equation with Wentzell boundary conditions and mass conservation*, Discrete Contin. Dyn. Systems **22** (2008), 1041-1063.
- [186] P. Galenko, D. Danilov and V. Lebedev, *Phase-field-crystal and Swift-Hohenberg equations with fast dynamics*, Phys. Rev. E **79** (2009), 051110.
- [187] P. Galenko and D. Jou, *Kinetic contribution to the fast spinodal decomposition controlled by diffusion*, Phys. A **388** (2009), 3113-3123.
- [188] P. Galenko and V. Lebedev, *Analysis of the dispersion relation in spinodal decomposition of a binary system*, Philos. Mag. Lett. **87** (2007), 821-827.
- [189] P. Galenko and V. Lebedev, *Local nonequilibrium effect on spinodal decomposition in a binary system*, Int. J. Thermodyn. **11** (2008), 21-28.
- [190] P. Galenko and V. Lebedev, *Nonequilibrium effects in spinodal decomposition of a binary system*, Phys. Lett. A **372** (2008), 985-989.
- [191] W. Gao and J. Yin, *System of Cahn-Hilliard equations with nonconstant interaction matrix*, Chinese Ann. Math., Ser. A **20** (1999), 169-176.
- [192] H. Garcke, *On Cahn-Hilliard systems with elasticity*, Proc. Roy. Soc. Edinburgh A **133** (2003), 307-331.
- [193] H. Garcke, *On a Cahn-Hilliard model for phase separation with elastic misfit*, Ann. Inst. H. Poincaré Anal. Non Linéaire **22** (2005), 165-185.
- [194] H. Garcke, *Mechanical effects in the Cahn-Hilliard model: A review on mathematical results*, in Mathematical methods and models in phase transitions, A. Miranville ed., Nova Sci. Publ., New York, 43-77, 2005.
- [195] H. Garcke, K.F. Lam, R. Nurnberg and E. Sitka, *A multiphase Cahn-Hilliard-Darcy model for tumour growth with necrosis*, Math. Models Methods Appl. Sci. **28** (2018), 525-577.
- [196] H. Garcke, K.F. Lam, E. Sitka and V. Styles, *A Cahn-Hilliard-Darcy model for tumour growth with chemotaxis and active transport*, Math. Models Methods Appl. Sci. **26** (2016), 1095-1148.
- [197] H. Garcke, K.F. Lam, E. Sitka and V. Styles, *Cahn-Hilliard inpainting with the double obstacle potential*, SIAM J. Imaging Sci. **11** (2018), 2064-2089.
- [198] H. Garcke and P. Knopf, *Weak solutions of the Cahn-Hilliard system with dynamic boundary conditions: A gradient flow approach*, preprint.
- [199] H. Garcke and U. Weikard, *Numerical approximation of the Cahn-Larché equation*, Numer. Math. **100** (2005), 639-662.
- [200] S. Gatti, M. Grasselli, A. Miranville and V. Pata, *On the hyperbolic relaxation of the one-dimensional Cahn-Hilliard equation*, J. Math. Anal. Appl. **312** (2005), 230-247.

- [201] S. Gatti, M. Grasselli, A. Miranville and V. Pata, *Hyperbolic relaxation of the viscous Cahn-Hilliard equation in 3-D*, Math. Models Methods Appl. Sci. **15** (2005), 165-198.
- [202] S. Gatti, M. Grasselli, A. Miranville and V. Pata, *Memory relaxation of first order evolution equations*, Nonlinearity **18** (2005), 1859-1883.
- [203] S. Gatti, M. Grasselli, A. Miranville and V. Pata, *Memory relaxation of the one-dimensional Cahn-Hilliard equation*, in Dissipative phase transitions, Ser. Adv. Math. Appl. Sci., Vol. **71**, World Sci. Publ., Hackensack, NJ, 101-114, 2006.
- [204] G. Giacomin and J.L. Lebowitz, *Phase segregation dynamics in particle systems with long range interaction I. Macroscopic limits*, J. Statist. Phys. **87** (1997), 37-61.
- [205] G. Giacomin and J.L. Lebowitz, *Phase segregation dynamics in particle systems with long range interaction II. Interface motion*, SIAM J. Appl. Math. **58** (1998), 1707-1729.
- [206] G. Gilardi, A. Miranville and G. Schimperna, *On the Cahn-Hilliard equation with irregular potentials and dynamic boundary conditions*, Commun. Pure Appl. Anal. **8** (2009), 881-912.
- [207] G. Gilardi, A. Miranville and G. Schimperna, *Long time behavior of the Cahn-Hilliard equation with irregular potentials and dynamic boundary conditions*, Chinese Ann. Math., Ser. B **31** (2010), 679-712.
- [208] A. Giorgini, M. Grasselli and A. Miranville, *The Cahn-Hilliard-Oono equation with singular potential*, Math. Models Methods Appl. Sci. **27** (2017), 2485-2510.
- [209] A. Giorgini, M. Grasselli and H. Wu, *The Cahn-Hilliard-Hele-Shaw system with singular potential*, Ann. Inst. H. Poincaré Anal. Non Linéaire **35** (2018), 1079-1118.
- [210] A. Giorgini, A. Miranville and R. Temam, *Uniqueness and regularity results for the Navier-Stokes-Cahn-Hilliard-Oono system with singular potentials*, submitted.
- [211] G.R. Goldstein and A. Miranville, *A Cahn-Hilliard-Gurtin model with dynamic boundary conditions*, Discrete Contin. Dyn. Systems Ser. S **6** (2013), 387-400.
- [212] G.R. Goldstein, A. Miranville and G. Schimperna, *A Cahn-Hilliard model in a domain with non-permeable walls*, Phys. D **240** (2011), 754-766.
- [213] A.A. Golovin, A.A. Nepomnyashchy, S.H. Davis and M.A. Zaks, *Convective Cahn-Hilliard models: from coarsening to roughening*, Phys. Rev. Lett. **86** (2001), 1550-1553.
- [214] H. Gomez, V.M. Calo, Y. Basilevs and T.J.R. Hughes, *Isogeometric analysis of Cahn-Hilliard phase field model*, Comput. Methods Appl. Mech. Engrg. **197** (2008), 4333-4352.
- [215] G. Gompper and M. Kraus, *Ginzburg-Landau theory of ternary amphiphilic systems. I. Gaussian interface fluctuations*, Phys. Rev. E **47** (1993), 4289-4300.
- [216] G. Gompper and M. Kraus, *Ginzburg-Landau theory of ternary amphiphilic systems. II. Monte Carlo simulations*, Phys. Rev. E **47** (1993), 4301-4312.
- [217] L. Goudenège, *Stochastic Cahn-Hilliard equation with singular nonlinearity and reflection*, Stochastic Process. Appl. **119** (2009), 3516-3548.
- [218] L. Goudenège and L. Manca, *Asymptotic properties of stochastic Cahn-Hilliard equation with singular nonlinearity and degenerate noise*, Stochastic Process. Appl. **125** (2015), 3785-3800.
- [219] L. Goudenège, D. Martin and G. Vial, *High order finite element calculations for the deterministic Cahn-Hilliard equation*, J. Sci. Comput. **52** (2012), 294-321.
- [220] C. Gräser, R. Kornhuber and U. Sack, *Nonsmooth Schur-Newton methods for multicomponent Cahn-Hilliard systems*, IMA J. Numer. Anal. **35** (2015), 652-679.
- [221] M. Grasselli, A. Miranville, R. Rossi and G. Schimperna, *Analysis of the Cahn-Hilliard equation with a chemical potential dependent mobility*, Commun. Partial Diff. Eqns. **36** (2011), 1193-1238.

- [222] M. Grasselli and M. Pierre, *A splitting method for the Cahn-Hilliard equation with inertial term*, Math. Models Methods Appl. Sci. **20** (2010), 1-28.
- [223] M. Grasselli and M. Pierre, *Energy stable and convergent finite element schemes for the modified phase field crystal equation*, ESAIM Math. Model. Numer. Anal. **50** (2016), 1523-1560.
- [224] M. Grasselli, G. Schimperna, A. Segatti and S. Zelik, *On the 3D Cahn-Hilliard equation with inertial term*, J. Evol. Eqns. **9** (2009), 371-404.
- [225] M. Grasselli, G. Schimperna and S. Zelik, *On the 2D Cahn-Hilliard equation with inertial term*, Commun. Partial Diff. Eqns. **34** (2009), 137-170.
- [226] M. Grasselli, G. Schimperna and S. Zelik, *Trajectory and smooth attractors for Cahn-Hilliard equations with inertial term*, Nonlinearity **23** (2010), 707-737.
- [227] M. Grasselli and H. Wu, *Well-posedness and longtime behavior for the modified phase-field crystal equation*, Math. Models Methods Appl. Sci. **24** (2014), 2743-2783.
- [228] M. Grasselli and H. Wu, *Robust exponential attractors for the modified phase-field crystal equation*, Discrete Contin. Dyn. Systems **35** (2015), 2539-2564.
- [229] Z. Guan, V. Heinonen, J.S. Lowengrub, C. Wang and S.M. Wise, *An energy stable, hexagonal finite difference scheme for the 2D phase field crystal amplitude equations*, J. Comput. Phys. **321** (2016), 1026-1054.
- [230] Z. Guan, J.S. Lowengrub and C. Wang, *Convergence analysis for second order accurate convex splitting schemes for the periodic nonlocal Allen-Cahn and Cahn-Hilliard equations*, submitted.
- [231] Z. Guan, J.S. Lowengrub, C. Wang and S.M. Wise, *Second-order convex splitting schemes for periodic nonlocal Cahn-Hilliard and Allen-Cahn equations*, J. Comput. Phys. **277** (2014), 48-71.
- [232] Z. Guan, C. Wang and S.M. Wise, *A convergent convex splitting scheme for the periodic nonlocal Cahn-Hilliard equation*, Numer. Math. **128** (2014), 377-406.
- [233] B. Guo, G. Wang and S. Wang, *Well posedness for the stochastic Cahn-Hilliard equation driven by Lévy space-time white noise*, Diff. Int. Eqns. **22** (2009), 543-560.
- [234] J. Guo, C. Wang, S.M. Wise and X. Yue, *An  $H^2$  convergence of a second-order convex-splitting, finite difference scheme for the three-dimensional Cahn-Hilliard equation*, Comm. Math. Sci. **14** (2016), 489-515.
- [235] M. Gurtin, *Generalized Ginzburg-Landau and Cahn-Hilliard equations based on a microforce balance*, Phys. D **92** (1996), 178-192.
- [236] M. Gurtin, D. Polignone and J. Vinals, *Two-phase binary fluids and immiscible fluids described by an order parameter*, Math. Models Methods Appl. Sci. **6** (1996), 815-831.
- [237] D. Han, *A decoupled unconditionally stable numerical scheme for the Cahn-Hilliard-Hele-Shaw system*, J. Sci. Comput. **66** (2016), 1102-1121.
- [238] D. Han and X. Wang, *Decoupled energy-law preserving numerical schemes for the Cahn-Hilliard-Darcy system*, Numer. Methods Partial Diff. Eqns. **32** (2016), 936-954.
- [239] D. Han and X. Wang, *A second order in time, decoupled, unconditionally stable numerical scheme for the Cahn-Hilliard-Darcy system*, J. Sci. Comput. **77** (2018), 1185-1209.
- [240] D. Han, X. Wang and H. Wu, *Existence and uniqueness of global weak solutions to a Cahn-Hilliard-Stokes-Darcy system for two phase incompressible flows in karstic geometry*, J. Diff. Eqns. **257** (2014), 3887-3933.
- [241] L. He and Y. Liu, *A class of stable spectral methods for the Cahn-Hilliard equation*, J. Comput. Phys. **228** (2009), 5101-5110.

- [242] B.S. Hosseini, S. Turek, M. Moller and C. Palmes, *Isogeometric analysis of the Navier-Stokes-Cahn-Hilliard equations with application to incompressible two-phase flows*, J. Comput. Phys. **348** (2017), 171-194.
- [243] Z. Hu, S.M. Wise, C. Wang and J.S. Lowengrub, *Stable finite difference, nonlinear multigrid simulation of the phase field crystal equation*, J. Comput. Phys. **228** (2009), 5323-5339.
- [244] B.R. Hunt and V.Y. Kaloshin, *Regularity of embeddings of infinite-dimensional fractal sets into finite-dimensional spaces*, Nonlinearity **12** (1999), 1263-1275.
- [245] B.R. Hunt, T. Sauer and J.A. Yorke, *Prevalence: A translation-invariant "almost every" for infinite-dimensional spaces*, Bull. Amer. Math. Soc. **27** (1992), 217-238.
- [246] Y. Huo, X. Jiang, H. Zhang and Y. Yang, *Hydrodynamic effects on phase separation of binary mixtures with reversible chemical reaction*, J. Chem. Phys. **118** (2003), 9830-9837.
- [247] Y. Huo, H. Zhang and Y. Yang, *Effects of reversible chemical reaction on morphology and domain growth of phase separating binary mixtures with viscosity difference*, Macromolecular Theory Simul. **13** (2004), 280-289.
- [248] S. Injrou and M. Pierre, *Stable discretizations of the Cahn-Hilliard-Gurtin equations*, Discrete Contin. Dyn. Systems **22** (2008), 1065-1080.
- [249] S. Injrou and M. Pierre, *Error estimates for a finite element discretization of the Cahn-Hilliard-Gurtin equations*, Diff. Int. Eqns. **15** (2010), 1161-1192.
- [250] H. Israel, *Long time behavior of an Allen-Cahn type equation with a singular potential and dynamic boundary conditions*, J. Appl. Anal. Comput. **2** (2012), 29-56.
- [251] H. Israel, *Well-posedness and long time behavior of an Allen-Cahn type equation*, Commun. Pure Appl. Anal. **12** (2013), 2811-2827.
- [252] H. Israel, A. Miranville and M. Petcu, *Well-posedness and long time behavior of a perturbed Cahn-Hilliard system with regular potentials*, Asymptot. Anal. **84** (2013), 147-179.
- [253] H. Israel, A. Miranville and M. Petcu, *Numerical analysis of a Cahn-Hilliard type equation with dynamic boundary conditions*, Ric. Mat. **64** (2015), 25-50.
- [254] A. Iuorio and S. Melchionna, *Long-time behavior of a nonlocal Cahn-Hilliard equation with reaction*, Discrete Contin. Dynam. Systems **38** (2018), 3765-3788.
- [255] D. Jacquemin, *Calculation of two phase Navier-Stokes flows using phase-field modeling*, J. Comput. Phys. **155** (1999), 96-127.
- [256] A.D. Jagtap and A.S. Vasudeva Murthy, *Higher order spectral element scheme for two- and three-dimensional Cahn-Hilliard equation*, Int. J. Adv. Engrg. Sci. Appl. Math. **10** (2018), 79-89.
- [257] J. Jiang, H. Wu and S. Zheng, *Well-posedness and long-time behavior of a nonautonomous Cahn-Hilliard-Darcy system with mass source modeling tumor growth*, J. Diff. Eqns. **259** (2015), 3032-3077.
- [258] G. Karali, and A. Katsoulakis, *The role of multiple microscopic mechanisms in cluster interface evolution*, J. Diff. Eqns. **235** (2007), 418-438.
- [259] G. Karali and Y. Nagase, *On the existence of solution for a Cahn-Hilliard/Allen-Cahn equation*, Discrete Contin. Dyn. Systems Ser. S **7** (2014), 127-137.
- [260] G. Karali and T. Ricciardi, *On the convergence of a fourth order evolution equation to the Allen-Cahn equation*, Nonlinear Anal. **72** (2010), 4271-4281.
- [261] A. Katsoulakis and G. Vlachos, *From microscopic interactions to macroscopic laws of cluster evolution*, Phys. Rev. , **84** (2000), 1511-1514.
- [262] D. Kay, V. Styles and E. Süli, *Discontinuous Galerkin finite element approximation of the Cahn-Hilliard equation with convection*, SIAM J. Numer. Anal. **47** (2009), 2660-2685.

- [263] D. Kay, V. Styles and R. Welford, *Finite element approximation of a Cahn-Hilliard-Navier-Stokes system*, Interfaces Free Bound. **10** (2008), 15-43.
- [264] N. Kenmochi, M. Niezgodka and I. Pawlow, *Subdifferential operator approach to the Cahn-Hilliard equation with constraint*, J. Diff. Eqns. **117** (1995), 320-356.
- [265] R. Kenzler, F. Eurich, P. Maass, B. Rinn, J. Schropp, E. Bohl and W. Dieterich, *Phase separation in confined geometries: solving the Cahn-Hilliard equation with generic boundary conditions*, Comput. Phys. Commun. **133** (2001), 139-157.
- [266] E. Khain and L.M. Sander, *A generalized Cahn-Hilliard equation for biological applications*, Phys. Rev. E **77** (2008), 051129.
- [267] J. Kim, *Phase field computations for ternary fluid flows*, Comput. Methods Appl. Mech. Engrg. **196** (2007), 4779-4788.
- [268] J. Kim, *Three-dimensional numerical simulations of a phase-field model for anisotropic interfacial energy*, Commun. Korean Math. Soc. **22** (2007), 453-464.
- [269] J. Kim, *A numerical method for the Cahn-Hilliard equation with a variable mobility*, Commun. Nonlinear Sci. Numer. Simul. **12** (2007), 1560-1571.
- [270] J. Kim, *A generalized continuous surface tension force formulation for phase-field models for multi-component immiscible fluid flows*, Comput. Methods Appl. Mech. Engrg. **198** (2009), 37-40.
- [271] J. Kim and K. Kang, *A numerical method for the ternary Cahn-Hilliard system with a degenerate mobility*, Appl. Numer. Math **59** (2009), 1029-1042.
- [272] J. Kim and J. Lowengrub, *Phase field modeling and simulation of three-phase flows*, Interfaces Free Bound. **7** (2005), 435-466.
- [273] I. Klapper and J. Dockery, *Role of cohesion in the material description of biofilms*, Phys. Rev. E **74** (2006), 0319021-0319028.
- [274] R. Kobayashi, *Modelling and numerical simulations of dendritic crystal growth*, Phys. D **63** (1993), 410-423.
- [275] R.V. Kohn and F. Otto, *Upper bounds for coarsening rates*, Commun. Math. Phys. **229** (2002), 375-395.
- [276] M. Korzec, P. Nayar and P. Rybka, *Global weak solutions to a sixth order Cahn-Hilliard type equation*, SIAM J. Math. Anal. **44** (2012), 3369-3387.
- [277] M. Korzec and P. Rybka, *On a higher order convective Cahn-Hilliard type equation*, SIAM J. Appl. Math. **72** (2012), 1343-1360.
- [278] A. Kostianko and S. Zelik, *Inertial manifolds for the 3D Cahn-Hilliard equations with periodic boundary conditions*, Commun. Pure Appl. Anal. **14** (2015), 2069-2094.
- [279] K.F. Lam and H. Wu, *Thermodynamically consistent Navier-Stokes-Cahn-Hilliard models with mass transfer and chemotaxis*, Eur. J. Appl. Math. **29** (2018), 595-644.
- [280] J.S. Langer, *Theory of spinodal decomposition in alloys*, Ann. Phys. **65** (1975), 53-86.
- [281] N. Lecoq, H. Zapolsky and P. Galenko, *Evolution of the structure factor in a hyperbolic model of spinodal decomposition*, Eur. Phys. J. Special Topics **177** (2009), 165-175.
- [282] H.G. Lee and J. Kim, *A second-order accurate non-linear difference scheme for the N-component Cahn-Hilliard system*, Phys. A **387** (2008), 19-20.
- [283] S. Lee, C. Lee, H.G. Lee and J. Kim, *Comparison of different numerical schemes for the Cahn-Hilliard equation*, J. Korean Soc. Ind. Appl. Math. **17** (2013), 197-207.
- [284] S. Lee and J. Shin, *Energy stable compact scheme for Cahn-Hilliard equation with periodic boundary condition*, Comput. Math. Appl., to appear.



- [285] C. Li, R. Qin, J. Ming and Z. Wang, *A discontinuous Galerkin method for stochastic Cahn-Hilliard equations*, *Comput. Math. Appl.* **75** (2018), 2100-2114.
- [286] D. Li and Z. Qiao, *On second order semi-implicit Fourier spectral methods for 2D Cahn-Hilliard equations*, *J. Sci. Comput.* **70** (2017), 301-341.
- [287] D. Li and C. Zhong, *Global attractor for the Cahn-Hilliard system with fast growing nonlinearity*, *J. Diff. Eqns.* **149** (1998), 191-210.
- [288] Y. Li, J. Kim and N. Wang, *An unconditionally energy-stable second-order time-accurate scheme for the Cahn-Hilliard equation on surfaces*, *Commun. Nonlinear Sci. Numer. Simul.* **53** (2017), 213-227.
- [289] Y. Li, H.G. Lee, B. Xia and J. Kim, *A compact fourth-order finite difference scheme for the three-dimensional Cahn-Hilliard equation*, *Comput. Phys. Commun.* **200** (2016), 108-116.
- [290] J.-L. Lions, *Quelques remarques sur les équations différentielles opérationnelles du 1er ordre*, *Rend. Sem. Mat. Padova* **33** (1963), 213-225.
- [291] C. Liu, *Convective Cahn-Hilliard equation with degenerate mobility*, *Dyn. Contin. Discrete Impuls. Systems Ser. A Math. Anal.* **16** (2009), 15-25.
- [292] C. Liu and J. Shen, *A phase field model for the mixture of two incompressible fluids and its approximation by a Fourier-spectral method*, *Phys. D* **179** (2003), 211-228.
- [293] C. Liu and H. Tang, *Existence of periodic solution for a Cahn-Hilliard/Allen-Cahn equation in two space dimensions*, *Evol. Eqns. Control Theory* **6** (2017), 219-237.
- [294] C. Liu and H. Wu, *An energetic variational approach for the Cahn-Hilliard equation with dynamic boundary conditions: derivation and analysis*, preprint.
- [295] F. Liu and J. Shen, *Stabilized semi-implicit spectral deferred correction methods for Allen-Cahn and Cahn-Hilliard equations*, *Math. Methods Appl. Sci.* **38** (2015), 4564-4575.
- [296] Q.-X. Liu, A. Doelman, V. Rottschäfer, M. de Jager, P.M.J. Herman, M. Rietkerk and J. van de Koppel, *Phase separation explains a new class of self-organized spatial patterns in ecological systems*, *Proc. Nation. Acad. Sci.*, <http://www.pnas.org/cgi/doi/10.1073/pnas.1222339110>.
- [297] S.-O. Londen and H. Petzeltova, *Regularity and separation from potential barriers for the Cahn-Hilliard equation with singular potential*, *J. Evol. Eqns.* **18** (2018), 1381-1393.
- [298] J. Lowengrub, E. Titi and K. Zhao, *Analysis of a mixture model of tumor growth*, *Eur. J. Appl. Math.* **24** (2013), 691-734.
- [299] J. Lowengrub and L. Truskinovski, *Quasi-incompressible Cahn-Hilliard fluids and topological transitions*, *Proc. Royal Soc. London Ser. A* **454** (1998), 2617-2654.
- [300] T. Ma and S. Wang, *Cahn-Hilliard equations and phase transition dynamics for binary systems*, *Discrete Contin. Dyn. Systems* **11** (2009), 741-784.
- [301] S. Maier-Paape, K. Mischaikow and T. Wanner, *Rigorous numerics for the Cahn-Hilliard equation on the unit square*, *Rev. Mat. Complutense* **21** (2008), 351-426.
- [302] S. Maier-Paape and T. Wanner, *Spinodal decomposition for the Cahn-Hilliard equation in higher dimensions. Part I: Probability and wavelength estimate*, *Commun. Math. Phys.* **195** (1998), 435-464.
- [303] S. Maier-Paape and T. Wanner, *Spinodal decomposition for the Cahn-Hilliard equation in higher dimensions: Nonlinear dynamics*, *Arch. Ration. Mech. Anal.* **151** (2000), 187-219.
- [304] A. Makki and A. Miranville, *Well-posedness for one-dimensional anisotropic Cahn-Hilliard and Allen-Cahn systems*, *Electron. J. Diff. Eqns.* **2015** (2015), 1-15.
- [305] A. Makki and A. Miranville, *Existence of solutions for anisotropic Cahn-Hilliard and Allen-Cahn systems in higher space dimensions*, *Discrete Contin. Dyn. Systems Ser. S* **9** (2016), 759-775.

- [306] H. Matsuoka and K.-I. Nakamura, *A stable finite difference method for a Cahn-Hilliard type equation with long-range interaction*, Sci. Rep. Kanazawa Univ. **57** (2013), 13-34.
- [307] S. Melchionna, H. Ranetbauer and L. Trussardi, *From nonlocal to local Cahn-Hilliard equation*, preprint.
- [308] S. Mikhailov, M. Hildebrand and G. Ertl, *Nonequilibrium nanostructures in condensed reactive systems*, in Coherent Structures in Complex Systems (Sitges, 2000), Vol. **567**, 252-269, 2001.
- [309] M.K. Miller, J.M. Hyde, M.G. Hetherington, A. Cerezo, G.D.W. Smith and C.M. Elliott, *Spinodal decomposition in Fe-Cr alloys: Experimental study at the atomic level and comparison with computer models I. Introduction and methodology*, Acta Metal. Mater. **43** (1995), 3385-3401.
- [310] M.K. Miller, J.M. Hyde, M.G. Hetherington, A. Cerezo, G.D.W. Smith and C.M. Elliott, *Spinodal decomposition in Fe-Cr alloys: Experimental study at the atomic level and comparison with computer models II. Development of domain size and composition amplitude*, Acta Metal. Mater. **43** (1995), 3403-3413.
- [311] M.K. Miller, J.M. Hyde, M.G. Hetherington, A. Cerezo, G.D.W. Smith and C.M. Elliott, *Spinodal decomposition in Fe-Cr alloys: Experimental study at the atomic level and comparison with computer models III. Development of morphology*, Acta Metal. Mater. **43** (1995), 3415-3426.
- [312] R.M. Mininni, A. Miranville and S. Romanelli, *Higher-order Cahn-Hilliard equations with dynamic boundary conditions*, J. Math. Anal. Appl. **449** (2017), 1321-1339.
- [313] S. Minjeaud, *An unconditionally stable uncoupled scheme for a triphasic Cahn-Hilliard/Navier-Stokes model*, Numer. Methods Partial Diff. Eqns. **29** (2013), 584-618.
- [314] A. Miranville, *Some generalizations of the Cahn-Hilliard equation*, Asymptotic Anal. **22** (2000), 235-259.
- [315] A. Miranville, *Long-time behavior of some models of Cahn-Hilliard equations in deformable continua*, Nonlinear Anal. Real World Appl. **2** (2001), 273-304.
- [316] A. Miranville, *Consistent models of Cahn-Hilliard-Gurtin equations with Neumann boundary conditions*, Phys. D **158** (2001), 233-257.
- [317] A. Miranville, *Asymptotic behavior of the Cahn-Hilliard-Oono equation*, J. Appl. Anal. Comput. **1** (2011), 523-536.
- [318] A. Miranville, *Asymptotic behaviour of a generalized Cahn-Hilliard equation with a proliferation term*, Appl. Anal. **92** (2013), 1308-1321.
- [319] A. Miranville, *Asymptotic behavior of a sixth-order Cahn-Hilliard system*, Cent. Eur. J. Math. **12** (2014), 141-154.
- [320] A. Miranville, *Sixth-order Cahn-Hilliard equations with logarithmic nonlinear terms*, Appl. Anal. **94** (2015), 2133-2146.
- [321] A. Miranville, *Sixth-order Cahn-Hilliard systems with dynamic boundary conditions*, Math. Methods Appl. Sci. **38** (2015), 1127-1145.
- [322] A. Miranville, *A generalized Cahn-Hilliard equation with logarithmic potentials*, Continuous and distributed systems II, Stud. Systems Decis. Control, Springer, Vol. **30**, 137-148, 2015.
- [323] A. Miranville, *On the phase-field-crystal model with logarithmic nonlinear terms*, Rev. Real Acad. Sci. **110** (2016), 145-157.
- [324] A. Miranville, *The Cahn-Hilliard equation and some of its variants*, AIMS Math. **2** (2017), 479-544.
- [325] A. Miranville and A. Piétrus, *A new formulation of the Cahn-Hilliard equation*, Nonlinear Anal. Real World Appl. **7** (2006), 285-307.

- [326] A. Miranville and A. Rougirel, *Local and asymptotic analysis of the flow generated by the Cahn-Hilliard-Gurtin equations*, Z. Angew. Math. Phys. **57** (2006), 244-268.
- [327] A. Miranville, W. Saoud and R. Talhouk, *Asymptotic behavior of a model for order-disorder and phase separation*, Asymptot. Anal. **103** (2017), 57-76.
- [328] A. Miranville, W. Saoud and R. Talhouk, *On The Cahn-Hilliard/Allen-Cahn equations with singular potentials*, Discrete Cont. Dynam. Systems Ser. B, to appear.
- [329] A. Miranville and G. Schimperna, *Nonisothermal phase separation based on a microforce balance*, Discrete Contin. Dyn. Systems Ser. B **5** (2005), 753-768.
- [330] A. Miranville and G. Schimperna, *Generalized Cahn-Hilliard equations for multicomponent alloys*, Adv. Math. Sci. Appl. **19** (2009), 131-154.
- [331] A. Miranville and G. Schimperna, *On a doubly nonlinear Cahn-Hilliard-Gurtin system*, Discrete Contin. Dyn. Systems Ser. B **14** (2010), 675-697.
- [332] A. Miranville and R. Temam, *On the Cahn-Hilliard-Oono-Navier-Stokes equations with singular potentials*, Appl. Anal. **95** (2016), 2609-2624.
- [333] A. Miranville and S. Zelik, *Robust exponential attractors for Cahn-Hilliard type equations with singular potentials*, Math. Methods Appl. Sci. **27** (2004), 545-582.
- [334] A. Miranville and S. Zelik, *Exponential attractors for the Cahn-Hilliard equation with dynamic boundary conditions*, Math. Methods Appl. Sci. **28** (2005), 709-735.
- [335] A. Miranville and S. Zelik, *Attractors for dissipative partial differential equations in bounded and unbounded domains*, in Handbook of Differential Equations, Evolutionary Partial Differential Equations, Vol. **4**, C.M. Dafermos and M. Pokorný eds., Elsevier, Amsterdam, 103-200, 2008.
- [336] A. Miranville and S. Zelik, *Doubly nonlinear Cahn-Hilliard-Gurtin equations*, Hokkaido Math. J. **38** (2009), 315-360.
- [337] A. Miranville and S. Zelik, *The Cahn-Hilliard equation with singular potentials and dynamic boundary conditions*, Discrete Contin. Dyn. Systems **28** (2010), 275-310.
- [338] Z. Mu, Y. Gong, W. Cai and Y. Wang, *Efficient local energy dissipation preserving algorithms for the Cahn-Hilliard equation*, J. Comput. Phys. **374** (2018), 654-667.
- [339] F. Nabet, *Convergence of a finite-volume scheme for the Cahn-Hilliard equation with dynamic boundary conditions*, IMA J. Numer. Anal. **36** (2016), 1898-1942.
- [340] T. Nagai, T. Senba and K. Yoshida, *Application of the Trudinger-Moser inequality to a parabolic system of chemotaxis*, Funkcial. Ekvac. **40** (1997), 411-433.
- [341] B. Nicolaenko and B. Scheurer, *Low dimensional behaviour of the pattern formation equations*, in Trends and practice of nonlinear analysis, V. Lakshmikantham ed., North-Holland, 1985.
- [342] B. Nicolaenko, B. Scheurer and R. Temam, *Some global dynamical properties of a class of pattern formation equations*, Commun. Partial Diff. Eqns. **14** (1989), 245-297.
- [343] A. Novick-Cohen, *Energy methods for the Cahn-Hilliard equation*, Quart. Appl. Math. **46** (1988), 681-690.
- [344] A. Novick-Cohen, *On the viscous Cahn-Hilliard equation*, in Material instabilities in continuum and related problems, J.M. Ball ed., Oxford University Press, Oxford, 329-342, 1988.
- [345] A. Novick-Cohen, *The Cahn-Hilliard equation: Mathematical and modeling perspectives*, Adv. Math. Sci. Appl. **8** (1998), 965-985.
- [346] A. Novick-Cohen, *Triple-junction motion for an Allen-Cahn/Cahn-Hilliard system*, Phys. D **137** (2000), 1-24.

- [347] A. Novick-Cohen, *The Cahn-Hilliard equation*, in Handbook of Differential Equations, Evolutionary Partial Differential Equations, Vol. 4, C.M. Dafermos and M. Pokorný eds., Elsevier, Amsterdam, 201-228, 2008.
- [348] Y. Oono and S. Puri, *Computationally efficient modeling of ordering of quenched phases*, Phys. Rev. Lett. **58** (1987), 836-839.
- [349] A. Oron, S.H. Davis and S.G. Bankoff, *Long-scale evolution of thin liquid films*, Rev. Mod. Phys. **69** (1997), 931-980.
- [350] I. Pawlow and G. Schimperna, *On a Cahn-Hilliard model with nonlinear diffusion*, SIAM J. Math. Anal. **45** (2013), 31-63.
- [351] I. Pawlow and G. Schimperna, *A Cahn-Hilliard equation with singular diffusion*, J. Diff. Eqns. **254** (2013), 779-803.
- [352] I. Pawlow and W.M. Zajaczkowski, *Strong solvability of 3-D Cahn-Hilliard system in elastic solids*, Math. Methods Appl. Sci. **32** (2008), 879-914.
- [353] I. Pawlow and W.M. Zajaczkowski, *Weak solutions to 3-D Cahn-Hilliard system in elastic solids*, Topol. Methods Nonlinear Anal. **32** (2008), 347-377.
- [354] I. Pawlow and W.M. Zajaczkowski, *Global regular solutions to Cahn-Hilliard system coupled with viscoelasticity*, Math. Methods Appl. Sci. **32** (2009), 2197-2242.
- [355] I. Pawlow and W.M. Zajaczkowski, *Long time behaviour of a Cahn-Hilliard system coupled with viscoelasticity*, Ann. Polon. Math. **98** (2010), 1-21.
- [356] I. Pawlow and W. Zajaczkowski, *A sixth order Cahn-Hilliard type equation arising in oil-water-surfactant mixtures*, Commun. Pure Appl. Anal. **10** (2011), 1823-1847.
- [357] I. Pawlow and W. Zajaczkowski, *On a class of sixth order viscous Cahn-Hilliard type equations*, Discrete Contin. Dyn. Systems Ser. S **6** (2013), 517-546.
- [358] M. Pierre, *Habilitation thesis*, Université de Poitiers, 2011.
- [359] P. Podio-Guidugli, *Models of phase segregation and diffusion of atomic species on a lattice*, Ric. Mat. **55** (2006), 105-118.
- [360] J. Prüss, R. Racke and S. Zheng, *Maximal regularity and asymptotic behavior of solutions for the Cahn-Hilliard equation with dynamic boundary conditions*, Ann. Mat. Pura Appl. (4) **185** (2006), 627-648.
- [361] J. Prüss, V. Vergara and R. Zacher, *Well-posedness and long-time behaviour for the non-isothermal Cahn-Hilliard equation with memory*, Discrete Contin. Dyn. Systems **26** (2010), 625-647.
- [362] T. Qian, X.-P. Wang and P. Sheng, *A variational approach to moving contact line hydrodynamics*, J. Fluid Mech. **564** (2006), 333-360.
- [363] R. Racke and S. Zheng, *The Cahn-Hilliard equation with dynamic boundary conditions*, Adv. Diff. Eqns. **8** (2003), 83-110.
- [364] A. Rajagopal, P. Fischer, E. Kuhl and P. Steinmann, *Natural element analysis of the Cahn-Hilliard phase-field model*, Comput. Mech. **46** (2010), 471-493.
- [365] J.C. Robinson, *Linear embeddings of finite-dimensional subsets of Banach spaces into Euclidean spaces*, Nonlinearity **22** (2009), 711-728.
- [366] A. Roggensack and C. Kraus, *Existence of weak solutions for the Cahn-Hilliard reaction model including elastic effects and damage*, J. Partial Diff. Eqns. **30** (2017), 111-145.
- [367] R. Rossi, *On two classes of generalized viscous Cahn-Hilliard equations*, Commun. Pure Appl. Anal. **4** (2005), 405-430.

- [368] R. Rossi, *Global attractor for the weak solutions of a class of viscous Cahn-Hilliard equations*, in Dissipative phase transitions, Ser. Adv. Math. Appl. Sci., Vol. **71**, World Sci. Publ., Hackensack, NJ, 247-268, 2006.
- [369] A. Rougirel, *Convergence to steady state and attractors for doubly nonlinear equations*, J. Math. Anal. Appl. **339** (2008), 281-294.
- [370] P. Rybka and K.-H. Hoffmann, *Convergence of solutions to Cahn-Hilliard equation*, Commun. Partial Diff. Eqns. **24** (1999), 1055-1077.
- [371] A. Sariyidin-Filibelioglu, B. Karaszen and M. Uzunca, *Energy stable interior penalty discontinuous Galerkin finite element method for Cahn-Hilliard equation*, Int. J. Nonlinear Sci. Numer. Simul. **18** (2017), 303-314.
- [372] G. Savaré and A. Visintin, *Variational convergence of nonlinear diffusion equations: applications to concentrated capacity problems with change of phase*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **8** (1997), 49-89.
- [373] A. Savostianov and S. Zelik, *Global well-posedness and attractors for the hyperbolic Cahn-Hilliard-Oono equation in the whole space*, Math. Models Methods Appl. Sci. **26** (2016), 1357-1384.
- [374] A. Savostianov and S. Zelik, *Finite dimensionality of the attractor for the hyperbolic Cahn-Hilliard-Oono equation in  $R^3$* , Math. Methods Appl. Sci. **39** (2016), 1254-1267.
- [375] R. Scala and G. Schimperna, *On the viscous Cahn-Hilliard equation with singular potential and inertial term*, AIMS Math. **1** (2016), 64-76.
- [376] L. Scarpa, *On the stochastic Cahn-Hilliard equation with a singular double-well potential*, Nonlinear Anal. **171** (2018), 102-133.
- [377] G. Schimperna, *Weak solution to a phase-field transmission problem in a concentrated capacity*, Math. Methods Appl. Sci. **22** (1999), 1235-1254.
- [378] G. Schimperna, *Global attractor for Cahn-Hilliard equations with nonconstant mobility*, Nonlinearity **20** (2007), 2365-2387.
- [379] G. Schimperna and S. Zelik, *Existence of solutions and separation from singularities for a class of fourth order degenerate parabolic equations*, Trans. Amer. Math. Soc. **365** (2013), 3799-3829.
- [380] J. Shen and X. Yang, *Numerical approximations of Allen-Cahn and Cahn-Hilliard equations*, Discrete Contin. Dyn. Systems **28** (2010), 1669-1691.
- [381] J. Shen and X. Yang, *Energy stable schemes for Cahn-Hilliard phase-field model of two-phase incompressible flows*, Chinese Ann. Math., Ser. B **31** (2010), 743-758.
- [382] W. Shen and S. Zheng, *On the coupled Cahn-Hilliard equations*, Commun. Partial Diff. Eqns. **18** (1993), 701-727.
- [383] J. Simon, *Compact sets in the space  $L^p(0, T; B)$* , Ann. Mat. Pura Appl. **CXLVI** (1987), 65-96.
- [384] J. Simon, *On the identification  $H = H'$  in the Lions theorem and a related inaccuracy*, Ric. Mat. **59** (2010), 245-255.
- [385] H. Song and C.-W. Shu, *Unconditional energy stability analysis of a second order implicit-explicit local discontinuous Galerkin method for the Cahn-Hilliard equation*, J. Sci. Comput. **73** (2017), 1178-1203.
- [386] R.H. Stogner, G.F. Carey and B.T. Murray, *Approximation of Cahn-Hilliard diffuse interface models using parallel adaptive mesh refinement and coarsening with  $C^1$  elements*, Int. J. Numer. Methods Engrg. **76** (2008), 636-661.
- [387] J.E. Taylor, *Mean curvature and weighted mean curvature*, Acta Metall. Mater. **40** (1992), 1475-1495.

- [388] R. Temam, *Infinite-dimensional dynamical systems in mechanics and physics*, Second edition, Applied Mathematical Sciences, Vol. **68**, Springer-Verlag, New York, 1997.
- [389] R. Temam, *Navier-Stokes equations: theory and numerical analysis*, AMS Chelsea Publishing, Vol. **343**, American Mathematical Society, Providence, Rhode Island, 2001.
- [390] C. Templier, Thèse de 3ème cycle, Université de Poitiers.
- [391] U. Thiele and E. Knobloch, *Thin liquid films on a slightly inclined heated plate*, Phys. D **190** (2004), 213-248.
- [392] G. Tierra and F. Guillén-González, *Numerical methods for solving the Cahn-Hilliard equation and its applicability to related energy-based models*, Arch. Comput. Methods Engrg. **22** (2015), 269-289.
- [393] S. Torabi, J. Lowengrub, A. Voigt and S. Wise, *A new phase-field model for strongly anisotropic systems*, Proc. R. Soc. A **465** (2009), 1337-1359.
- [394] S. Tremaine, *On the origin of irregular structure in Saturn's rings*, Astron. J. **125** (2003), 894-901.
- [395] J.D. van der Waals, *The thermodynamic theory of capillarity flow under the hypothesis of a continuous variation of density (in Dutch)*, Verhandel. Konink. Akad. Wetten Amsterdam (Sec. 1), **Vol. 1**, 1893.
- [396] J. Verdasca, P. Borckmans and G. Dewel, *Chemically frozen phase separation in an adsorbed layer*, Phys. Review E **52** (1995), 4616-4619.
- [397] S. Villain-Guillot, *Phases modulées et dynamique de Cahn-Hilliard*, Habilitation thesis, Université Bordeaux I, 2010.
- [398] C. Wang and S.M. Wise, *Global smooth solutions of the modified phase field crystal equation*, Methods Appl. Anal. **17** (2010), 191-212.
- [399] C. Wang and S.M. Wise, *An energy stable and convergent finite difference scheme for the modified phase field crystal equation*, SIAM J. Numer. Anal. **49** (2011), 945-969.
- [400] J. Wang, Q. Zhai, R. Zhang and S. Zhang, *A weak Galerkin finite element scheme for the Cahn-Hilliard equation*, Math. Comp. **88** (2019), 211-235.
- [401] W. Wang, L. Chen and J. Zhou, *Postprocessing mixed finite element methods for solving Cahn-Hilliard equation: methods and error analysis*, J. Sci. Comput. **67** (2016), 724-746.
- [402] X. Wang and H. Wu, *Long-time behavior for the Hele-Shaw-Cahn-Hilliard system*, Asymptot. Anal. **78** (2012), 217-245.
- [403] L. Wang and H. Yu, *Convergence analysis of an unconditionally energy stable linear Crank-Nicolson scheme for the Cahn-Hilliard equation*, J. Math. Study **51** (2018), 89-114.
- [404] X. Wang and Z. Zhang, *Well-posedness of the Hele-Shaw-Cahn-Hilliard system*, Ann. Inst. H. Poincaré Anal. Non Linéaire **30** (2013), 367-384.
- [405] S.J. Watson, F. Otto, B. Rubinstein and S.H. Davis, *Coarsening dynamics of the convective Cahn-Hilliard equation*, Phys. D **178** (2003), 127-148.
- [406] G.N. Wells, E. Kuhl and K. Garikipati, *A discontinuous Galerkin method for Cahn-Hilliard equation*, J. Comput. Phys. **218** (2006), 860-877.
- [407] A.A. Wheeler and G.B. McFadden, *On the notion of  $\xi$ -vector and stress tensor for a general class of anisotropic diffuse interface models*, Proc. R. Soc. London Ser. A **453** (1997), 1611-1630.
- [408] M. Wilke,  *$L_p$ -theory for a Cahn-Hilliard-Gurtin system*, Evol. Eqns. Control Theory **1** (2012), 393-429.
- [409] S.M. Wise, *Unconditionally stable finite difference, nonlinear multigrid simulations of the Cahn-Hilliard-Hele-Shaw system of equations*, J. Sci. Comput. **44** (2010), 38-68.

- [410] S. Wise, J. Kim and J. Lowengrub, *Solving the regularized, strongly anisotropic Cahn-Hilliard equation by an adaptive nonlinear multigrid method*, J. Comput. Phys. **226** (2007), 414-446.
- [411] S.M. Wise, C. Wang and J.S. Lowengrub, *An energy stable and convergent finite difference scheme for the phase field crystal equation*, SIAM J. Numer. Anal. **47** (2009), 2269-2288.
- [412] H. Wu and S. Zheng, *Convergence to equilibrium for the Cahn-Hilliard equation with dynamic boundary conditions*, J. Diff. Eqns. **204** (2004), 511-531.
- [413] Y. Xia, Y. Xu and C.-W. Shu, *Application of the local discontinuous Galerkin method for the Allen-Cahn/Cahn-Hilliard system*, Commun. Comput. Phys. **5** (2009), 821-835.
- [414] X. Yang and H. Yu, *Efficient second order unconditionally stable schemes for a phase field moving contact line model using an invariant energy quadratization approach*, SIAM J. Sci. Comput. **40** (2018), B889-B914.
- [415] X. Yang and J. Zhao, *Efficient linear schemes for the nonlocal Cahn-Hilliard equation of phase field models*, Comput. Phys. Commun., to appear.
- [416] X. Yang, J. Zhao and X. He, *Linear, second order and unconditionally energy stable schemes for the viscous Cahn-Hilliard equation with hyperbolic relaxation using the invariant energy quadratization method*, J. Comput. Appl. Math. **343** (2018), 80-97.
- [417] J. Yin and C. Liu, *Cahn-Hilliard type equations with concentration dependent mobility*, in Mathematical methods and models in phase transitions, A. Miranville ed., Nova Sci. Publ., New York, 79-93, 2005.
- [418] B. You and F. Li, *Well-posedness and global attractor of the Cahn-Hilliard-Brinkman system with dynamic boundary conditions*, Dyn. Partial Diff. Eqns. **13** (2016), 75-90.
- [419] P. Yue, J.J. Feng, C. Liu and J. Shen, *A diffuse-interface method for simulating two-phase flows of complex fluids*, J. Fluid Mech. **515** (2004), 293-317.
- [420] S. Zelik, *The attractor for a nonlinear reaction-diffusion system with a supercritical nonlinearity and its dimension*, Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. **24** (2000), 1-25.
- [421] T. Zhang and Q. Wang, *Cahn-Hilliard vs singular Cahn-Hilliard equations in phase field modeling*, Commun. Comput. Phys. **7** (2010), 362-382.
- [422] X. Zhang and C. Liu, *Existence of solutions to the Cahn-Hilliard/Allen-Cahn equation with degenerate mobility*, Electron. J. Diff. Eqns. **2016** (2016), 1-22.
- [423] L.-Y. Zhao, H. Wu and H.-Y. Huang, *Convergence to equilibrium for a phase-field model for the mixture of two incompressible fluids*, Commun. Math. Sci. **7** (2009), 939-962.
- [424] S. Zheng, *Asymptotic behavior of solution to the Cahn-Hilliard equation*, Appl. Anal. **23** (1986), 165-184.
- [425] X. Zheng, C. Yang, X.-C. Cai and D. Keyes, *A parallel domain decomposition-based implicit method for the Cahn-Hilliard-Cook phase-field equation in 3D*, J. Comput. Phys. **285** (2015), 55-70.
- [426] J.X. Zhou and M.E. Li, *Solving phase field equations with a meshless method*, Commun. Numer. Methods Engrg. **22** (2006), 1109-1115.

UNIVERSITÉ DE POITIERS, POITIERS, FRANCE  
FUDAN UNIVERSITY (FUDAN FELLOW), SHANGHAI, CHINA  
XIAMEN UNIVERSITY, XIAMEN, FUJIAN, CHINA  
Email address: Alain.Miranville@math.univ-poitiers.fr